

Nest Population Size and Potential Production of Geese and Spectacled Eiders on the Yukon-Kuskokwim Delta, Alaska, 2008



Julian B. Fischer¹, Robert A. Stehn¹, and George Walters²

¹ U.S. Fish and Wildlife Service, Migratory Bird Management, Waterfowl Management Branch, 1011 E. Tudor Rd., Anchorage, AK 99503

² U.S. Fish and Wildlife Service, Yukon Delta National Wildlife Refuge, P.O. Box 346, Bethel, AK 99559

SUMMARY: A ground-based survey has been used since 1985 to estimate the number of nests and eggs of geese and eiders on the Yukon-Kuskokwim Delta (YKD), Alaska. 2008 was a year of fair to good production for geese and eiders. Depredation and nest abandonment rates were moderate for most species. Recent 10-year trends in nests populations are stable for all waterfowl except for spectacled and common eiders and white-fronted geese that are continuing to grow significantly. We estimated that spectacled eiders built 4,991 nests and laid 19,321 viable eggs on the YKD coastal zone in 2008. The estimate of spectacled eider nests was 8% above the 1985-2007 average. This year marked the first time in the history of the survey that the recent 10-year average growth rate for spectacled eiders is significantly above 1.00 ($\alpha > 0.90$). Spring arrived later on the YKD in 2008 compared to recent years, but a long-term advance in hatch dates is apparent over the course of this survey.

INTRODUCTION:

Annual assessment of nest population size and egg production of geese and eiders on the Yukon-Kuskokwim Delta (YKD) provides information for the Spectacled and Steller's Eider Recovery Team, the Pacific Flyway Technical Committee, participants in cooperative goose management plans, and biologists interested in waterfowl status and trends in western Alaska. A ground-based sampling procedure has been used since 1985 to estimate the number of nests and eggs for spectacled eiders (*Somateria fischeri*), cackling geese (*Branta hutchinsii minima*), emperor geese (*Chen canagica*), greater white-fronted geese (*Anser albifrons frontalis*), and other nesting waterbirds on the YKD. The ground-based nest survey provides an estimate of nest population size and potential production. The ground survey is conducted concurrently with an aerial breeding pair survey (Bollinger and Eldridge 2008, Platte and Stehn 2008) that provides an index to population size. Together, these surveys contribute long-term data needed to understand goose and eider population status and reproductive success.



Starting in 2006 we improved analysis methods by incorporating nest detection rates (Bowman and Stehn *manuscript in prep.*), reducing sampling error in expansion factors, and restricting analyses of historical data to plots within the current survey area. These changes resulted in population estimates that differ from those reported in previous annual reports (Fischer et al. 2007).

METHODS:

We used a ground-based sampling procedure to monitor goose and eider nest populations and potential production on the YKD coastal zone from 1985 to 2008 (Fig. 1). Boundaries of the survey area included all lands on the Yukon Delta National



Wildlife Refuge (YDNWR) containing medium and high nest densities of spectacled eiders (based on aerial and ground observations 1985-1993, USFWS unpubl. data). We excluded privately owned high density nesting habitat near Kokechik Bay, two patches on south Nelson Island, and several tracts near Hazen Bay because annual access could not be assured. From 1994-1997 and 2000-2008, plots were selected within 716 km² (core spectacled eider habitat; hereafter “core”), which comprised 5.6% of the total coastal zone. In 1998, 1999, and prior to 1994, plots were selected within a slightly a larger portion of the coastal zone. In this report, estimates of nest population size and egg production are based on plots within the core

716 km², whereas estimates of clutch size and hatch date use data from plots within and beyond the core area.

We used GIS and custom-written TrueBASIC computer programs to randomly select 85 plots within the core 716 km² ground sampled area in 2008 (Fig. 1). Areas sampled during the preceding five years were excluded from the random selection process. We transferred plot boundaries to digital color infrared aerial photographs (1:15,000 or 1:10,000) and IKONOS satellite imagery for field use. We included plots regardless of juxtaposition to lakes and rivers. Plot size was 402 m by 805 m (0.32 km²) in 1988-1994 and 1997-2008. Plot sizes were variable in 1985-1987 (0.16-1.66 km²), 0.45 km² in 1995, and 0.36 km² in 1996.

Plots were searched by 2-4 biologists who were transported either by Cessna 185 float-equipped aircraft or by motorboat. Two boat crews originated from the YDNWR Kanaryamiut Field Station and worked plots accessible from the Aphrewn and Opagarak rivers. A third boat crew was transported directly from Bethel to plots on the Naskonat Peninsula, directly north of Kigigak Island. Biologists at Kigigak Island (YDNWR), Big Slough (USGS), Manokinak River (USGS), and Tutakoke River (University of Nevada, Reno) searched plots near their camps. All sites dry enough for a nest were examined for active and destroyed waterfowl, crane, loon, and gull nests. Incidental nesting records of other species were recorded as encountered (data available upon request), but most shorebird and passerine nests were likely missed.




Figure 1. Location of 85 randomly selected plots within the ground sampled area (716 km²) relative to the Yukon-Kuskokwim Delta coastal zone (12,832 km²), Alaska, 2008. Sampled plots are represented by solid rectangles. Three additional plots were selected but not sampled in 2008.

At each nest we recorded species, nest status, nest site, and stage of incubation. Species was determined by visual confirmation of an adult at the nest or by comparing down and contour feathers in the nest bowl with a photographic field guide (Bowman 2008). We determined stage of incubation for all species by measuring float angles of eggs from active nests (Westerskov 1950). Hatch dates were estimated from incubation stage of eggs found on plots. Hatch date estimates prior to 1985 were derived from plots established by Butler (1983).

Red-throated (*Gavia stellata*) and Pacific loons (*Gavia pacifica*) rarely remain near their nest sites when disturbed by plot searchers, and their nests are essentially indistinguishable from each other (Bowman 2008), so we applied the ratio of aerial observations of each loon species, as recorded from an aerial platform (Platte and Stehn 2008) to determine the relative numbers of loon nests for each species. To reduce variation in the ratio of loon species caused by sampling error in aerial indices, we used a localized 7-year average (from the current year, 3 prior years, and 3 following years, as available) of the annual ratio of loon species. For example, the localized 7-year average ratio for 1995 was the mean of ratios from 1992-1998; whereas for 2008 we use the mean of ratios from years 2005 to present.



The mean and variance of the number of nests and eggs per plot was based on a simple random sample of plots. The estimates of nests and eggs were expanded to the ground sampled area (716 km²) based on the size and number of plots for each year. Nest population estimates were corrected for detection rate using a model that considers species, nest activity status, observer experience, and nest site location (Bowman and Stehn *manuscript in prep.*).

The corrected estimates for nests and eggs in the ground sampled area were expanded to the entire coastal zone of the YKD based on a stratified analysis of an aerial survey of the entire YKD coastal zone (12,832 km²; Butler et al. 1988, Bollinger and Eldridge 2008, Platte and Stehn 2008). To expand the ground-based estimates, we annually calculated the ratio of the aerial breeding population index outside the ground-sampled area (“OUT”) to the aerial index within the ground-sampled area (“IN”). We used the localized 7-year average (from the current year, 3 prior years, and 3 following years, as available) of 1+ Out/In ratio as an expansion factor for a given year. Variance estimates of nest populations expanded to the entire coastal zone incorporate the variance of the Out/In ratio. The aerial breeding population index for most species was based on twice the number of singles plus the number of birds in pairs observed, because single geese, cranes, and ducks observed are assumed to be the mates of unobserved females on nests. Flocks of these species were not included in the aerial index, except with brant. For swans, the number of single birds observed was not doubled. For loons and gulls, the total number of birds observed was used as the index. The expansion of estimated nests and eggs from the ground sampled area to the entire coastal zone is based the assumption that observed breeding bird indices obtained from aerial surveys provide an accurate linear relationship with the number of nests within versus outside of the ground sampled area.

Data were tabulated, edited, sorted and analyzed using Excel, and nest population, hatch date and clutch size estimates were calculated using customized TrueBASIC programs.

The estimated total number of nests is a direct measure of effective breeding population size and an index to the size of the population of adults that are potential nesters. The estimated total number of eggs is a measure of the number of young that could potentially augment the fall population if they survive through summer. The proportion of nests that are active when the plots are searched is an index to nest success; the actual proportion of nests that produced young is lower because some nests are lost after plots are searched. The relative production rate measures how the number of eggs per total nests constructed in a given year compares to the long-term average for each species; thereby quantifying the relative annual contribution of eggs to the population on a per pair basis.



RESULTS:

We searched 82 plots from 1 June to 18 June (Fig. 1). Crews based at the Kanaryamiut Field Station searched 35 plots, boat-based crews searched 33 plots, a Yukon Delta NWR crew at Kigigak Island searched three plots, a USGS crew at Big Slough searched four plots, a USGS crew at the Manokinak River searched four plots, and a University of Nevada, Reno crew at Tutakoke River searched three plots. Three of the 85 randomly selected plots were not sampled due to access and weather delays that prevented completion of field work prior to onset of hatch. Crews located 3,694 nests within plot boundaries in 2008 comprised of 1,579 cackling goose, 355 emperor goose, 600 greater white-fronted goose, 108 spectacled eider, and 1,052 nests of other species. Calculations of clutch size and hatch date also included an additional 43 nests located outside of plot boundaries. We present nest population, egg production, and nest success estimates in figures with accompanying tabulated data for each species (Fig. 2). Estimated hatch date for each species is presented in Table 1. The following section presents general descriptive results for each species.

Cackling Geese (*Branta hutchinsii minima*)

Production of cackling geese was fair in 2008. Total numbers of nests were down from 2007 (Fig. 2) but remained above the long-term (1985-2007) and short-term (1998-2007) averages. Egg production in 2008 was higher than the long-term average; however, owing to relatively low clutch size and nest success, the overall egg production was down from 2007 and below the short-term mean. As a result, the relative production rate of eggs per total nest



(i.e. number of breeding pairs) was below average. Following thirteen years of positive

growth, the growth rate of cackling goose nests and eggs has been essentially flat during the last decade. The estimates of number of nests and eggs were fifth and sixth highest, respectively, since 1985. Average hatch date for cacklers in 2008 was three days later than 2007 and two days earlier than the previous long-term average (1985-2007; Table 1).

Emperor Geese (*Chen canagica*)



Potential production of emperor geese was good in 2008. Total numbers of nests and eggs were down from 2007 (Fig. 2) but remained above the long-term (1985-2007) and short-term (1998-2007) averages. Nest success and clutch size were similar to short and long-term means, and resulted in a relative production rate just below the long-term mean. Nest and egg population size have been variable over the course of this survey and have resulted in non-significant growth rates in both short- and long-term time spans.

Similar to other waterfowl species, estimated hatch date for emperors in 2008 was two days later than 2007 and one day earlier than the previous long-term average (1985-2007; Table 1).

Greater White-fronted Geese (*Anser albifrons frontalis*)

Potential production of white-fronted geese was poor to fair in 2008. Total numbers of nests and eggs were down from 2007 (Fig. 2) but remained above the long-term (1985-2007) and short-term (1998-2007) averages. In contrast, egg production in 2008 was down sharply from the previous four years (Fig. 2) resulting in lower than average production in the short term (1998-2007).

Clutch size and nest success were both lower than short and long-term averages. As a result, the relative production rate in 2008 was below average, and the third lowest value in 24 years. White-front nest and egg population growth rates have been high throughout the duration of the survey but



have slowed somewhat during the last decade. Similar to other waterfowl species, estimated hatch date for white-fronts in 2008 was three days later than 2007 and one day earlier than the previous long-term average (1985-2007; Table 1).

Black Brant (*Branta bernicla nigricans*)

The nest plot survey was not designed to monitor colonial nesting birds such as black brant, and we purposely excluded most primary colonies from the sample area. Kigigak Island supports one of the primary brant colonies on the YKD and is included in the ground sampled area. Five primary brant colonies are monitored annually by digital photographic surveys (see Wilson 2008). The nest plot survey does provide an estimate of nests populations from non-colonial brant, small satellite colonies, and one of the five major YKD colonies (Kigigak Island). In these areas, potential production of brant in

2008 was fair to good. The number of nests in 2008 was down from the previous year. Nest population estimates were slightly below the long-term average, and slightly above the short-term average. Clutch size in 2008 was higher than the short-term mean, but lower than the long-term mean, whereas nest success was higher than short and long-term averages. As a result, the relative production rate in 2008 was very close to average. Nest and egg population sizes have been variable over the course of this survey and resulted in non-significant growth rates in both short- and long-term time spans. Similar to other waterfowl species, estimated hatch date for brant in 2008 was two days later than 2007 and one day earlier than the previous long-term average (1985-2007; Table 1).

Tundra Swans (*Cygnus columbianus*)

Potential production of tundra swans was fair in 2008. Total numbers of nests and eggs were down from 2007 (Fig. 2) and were below short- (1998-2007) and long-term (1985-2007) averages. Clutch size was also down from 2007 and below short- and long-term averages. Nest success in 2008, was unchanged from the short-term average but down from the long-term mean. As a result, the relative production rate in 2008 was below average. For tundra swans, nest and egg population sizes have been variable over the course of this survey and resulted in non-significant growth rates in the short-term. Over the long-term, however, both nest and egg population sizes have increased significantly. Unlike other waterfowl species, estimated hatch date for tundra swans in 2008 was later than 2007 by three days, and one day later than the previous long-term average (1985-2007; Table 1).

Sandhill Cranes (*Grus canadensis*)



Potential production of sandhill cranes was good in 2008. Total numbers of nests and eggs were down from 2007 (Fig. 2) and were below the short- (1998-2007) and long-term (1985-2007) averages. Similarly, clutch size in 2008 was below short and long-term means. Despite these declines, nest success of those cranes that bred was higher than short- and long-term averages resulting in an average relative production rate. Nest and egg population sizes have been variable over the course of this survey and have resulted in non-significant growth rates in both short- and long-term time spans. Unlike the waterfowl species, estimated hatch date for sandhill cranes in 2008 was over a week later than 2007 and one day later than the previous long-term average (1985-2007; Table 1).

Spectacled Eiders (*Somateria fischeri*)

Potential production of spectacled eiders was good in 2008. Total numbers of nests and eggs were up from 2007 (Fig. 2) and remained above the short-term (1998-2007) and long-term (1985-2007) averages. Clutch size in 2008 was higher than short- and long-term means. However, nest success was down sharply from 2007 and was below short- and long-term averages. Together, these measures resulted in



an average relative production rate. Estimated numbers of nests have grown since the early 1990s, and this change is reflected in a significant positive short-term growth rate (1999-2008). In contrast, the long-term growth rate (1985-2008) is negative, largely due to the high estimates measured in the mid and late 1980s. Annual estimates of egg production are variable due to fluctuations in breeding conditions and have led to non-significant trends in short- and long-term time periods. Estimated hatch date for spectacled eiders in 2008 was the same as in 2007 and two days earlier than the previous long-term average (1985-2007; Table 1).

Common Eiders (*Somateria mollissima*)

Potential production of common eiders was good in 2008. Total numbers of nests and eggs were down sharply from 2007 (Fig. 2) but remained above the short-term (1998-2007) and long-term (1985-2007) averages. Clutch size in 2008 was very similar to short- and long-term means. Nest success was down from 2007 but remained above the short- and long-term average. Together, these measures resulted in an average relative production rate. Nest numbers and egg production have grown throughout the course of this survey. This change is reflected in significant positive short- (1999-2008) and long-term (1985-2008) growth rates. Similar to other waterfowl species, estimated hatch date for common eiders in 2008 was four days later than 2007 and two days earlier than the previous long-term average (1985-2007; Table 1).



Loons

Potential production of Pacific loons (*Gavia pacifica*) and red-throated loons (*Gavia stellata*) was fair in 2008. Estimates of nests and eggs for both species were close to estimates in 2007 and remained above the short- and long-term averages. While nest success for both species in 2008 was similar to short- and long-term means, clutch sizes were down sharply, resulting in a lower than average relative production rate. Nest and egg population sizes have been variable over the course of this survey and resulted in non-significant growth rates in both short- and long-term time spans. Similar to waterfowl species, estimated hatch date for both loon species in 2008 was four days later than 2007 and two days earlier than the previous long-term average (1985-2007; Table 1).

Gulls and Terns



Colonial nesting species such as gulls (glaucous gulls [*Larus hyperboreus*], Sabine's gulls [*Xema sabini*], mew gulls [*Larus canus*]), and arctic terns [*Sterna paradisaea*) are not monitored with precision by the nest plot survey. Nonetheless, the survey does provide a measure of potential production for these species. In 2008, potential production was good for gulls and terns (Fig. 2). Numbers of nests and eggs in 2008 was down from the previous year for all but mew gulls, but estimates were higher than the

short- and long-term averages. Clutch size for all three gull species was close to short- and long-term averages. Nest success for most gulls was higher than short- and long-term averages. The exception was mew gulls whose nest success was relatively low. In contrast, terns had lower than average clutch size and higher than average nest success in 2008. Trends in nest populations of gulls and terns are highly variable over the course of the survey; however, positive growth in the nest populations of mew gulls is apparent during the most recent decade. Estimated hatch date of gulls and terns in 2008 was up to five days later than 2007 and up to four days earlier than the long-term average (Table 1).

DISCUSSION:

The nest plot survey was designed to provide annual estimates of nest population size and trend, and predict potential production (active eggs) and hatching dates for nesting geese (cackling geese, emperor geese, greater white-fronted geese) and eiders (spectacled and common) on the Yukon-Kuskokwim Delta coastal zone.

In general, production in 2008 was fair for cackling and white-fronted geese, and good for emperor geese and eiders. We observed low nest success for cackling and white-fronted geese and spectacled eiders, but overall numbers of nests with eggs helped offset low success. Among the principal species monitored in the survey, only common eiders exhibited relatively high nest success. The relative production rate (contribution of eggs per bird relative to the long-term average) was below the median for all species except glaucous gulls. For white-fronted geese, 2008 represented the third lowest relative production rate during the 24 years of data collection.

Incidence of nest desertion and depredation was variable among plots and not isolated to one region within the ground sampled area. Rather, nest failure was distributed throughout the study area, and plots with severe nest loss were juxtaposed with plots with high nest success. Arctic fox (*Alopex lagopus*) have been identified as important nest depredators on the coastal zone, particularly in brant colonies (Anthony et al. 1991). While fox were likely contributors to egg loss on study plots in 2008, there is no indication that fox presence necessarily resulted in high levels of depredation in all instances. For example, waterfowl nest success on plots with fox sign (fur, tracks, active dens, fox observed) ranged from 50% to 100%. Further, mean nest success was virtually the same among plots with and without sign of fox presence (84% and 86% success, respectively). Most primary brant colonies were excluded from the study area; thus we do not have data to measure the degree of fox predation within those colonies.

Spring was cold and relatively late in 2008. The Kuskokwim River broke up at Bethel on May 13, ten days later than 2007 and two days later than the previous 26-year average (NOAA 2008). Timing of waterfowl nest initiation is closely correlated with spring breakup (Raveling 1978, Dau and Mickelson 1979). Consequently, in 2008 waterfowl nests hatched approximately four days later than the previous year and two days earlier than the long-term averages. A significant trend in earlier waterfowl nest initiation is apparent over the 27-year history of this study (Fig. 3). Based on nesting data since 1982, we estimate that on average, hatch for cackling geese for which the most data are available, occurs 0.4 days earlier each year. This change translates to an advance of over 10 days during the 1982-2008 time period.

In general, goose nest population trends parallel trends derived from aerial breeding pair surveys (Bollinger and Eldridge 2008, Platte and Stehn 2008). Estimates of cackling goose nests were at record lows in the mid-1980s prior to adoption of the

cooperative Yukon-Kuskokwim Delta Goose Management Plan that provided much needed protection for nesting and wintering populations of geese (Pamplin 1986). Data from this nest plot survey and aerial breeding pair surveys (Bollinger and Eldridge 2008, Platte and Stehn 2008) show that by the late-1980s, the cackling goose population of nests and pairs increased rapidly, peaking in the late-1990s. Since 1999, trends for



cackling geese have been relatively flat (this study, Bollinger and Eldridge 2008). Unlike the other goose species, populations of emperor geese did not increase markedly after adoption of the Yukon Delta Goose Management Plan, although a slow annual increase in the long-term trends from ground and air surveys is apparent (this study, Bollinger and Eldridge 2008). The population of greater white-

fronted geese has increased dramatically on the coastal zone of the YKD since the mid-1980s and has continued to grow at moderate levels in recent years (this study, Bollinger and Eldridge 2008).

The number of spectacled eider nests has varied widely since the beginning of the survey in the mid-1980s. The average long-term growth rate (1985-2008) is negative, but the rate is not significantly different from 1.00. However, for the first time in the history of this study, the recent 10-year growth rate is significantly above 1.00 ($\alpha > 0.90$).

Aerial surveys of spectacled eider pairs on the YKD show a similar contemporary increasing trend as the nest population size trend (Platte and Stehn 2008). The aerial survey indices are influenced by survey timing, and an analysis of aerial data from 1993-2006 that corrected for survey timing reported a growth rate of 1.042 (Stehn et al. 2006), a rate nearly identical to the nest population trend from the most recent 10 years.

Spectacled eider nest success is variable among years and generally reflects varying levels of nest predation. Plots are visited one time, so the measure of nest success is an overestimation of actual success because some nests undoubtedly fail after the plot is searched. Nonetheless, the pattern in nest success measured from the nest plot survey (number of active nests divided by total nests times 100%, corrected for detection rate) generally matches nest success at Kigigak Island (successful hatched nests/total nests) where nests are visited every seven days until hatch (Lake 2008). In 2008, the nest plot survey and intensive surveys at Kigigak both reported relatively low nest success (Fig. 4). The largest difference between these measures of nest success from these two surveys was noted in 2001 and 2003, years of very poor production, where perhaps many nest failures occurred late in nesting. Alternatively, a localized factor may have caused low success on Kigigak Island during those years. Large annual fluctuations in production estimates are reduced when sampling occurs over a large portion of the nesting range, such as in the nest plot survey described here.

A comparison of estimates from the photographic colony survey and the ground plot survey suggests that many brant occur as dispersed nesters or in satellite colonies outside most primary colonies (Fig. 5). Photographic methods were initiated in the 1990s to monitor the nest population of brant in major colonies with greater precision than ground surveys or standard straight line aerial waterfowl surveys (Wilson 2008). Estimates of brant nests derived from ground plot and photographic surveys both indicate a slow decline in nest populations over the last 17 years (Fig. 5).

The population sizes of goose and eider nests should not be interpreted as direct estimates of population size. For example, a year of poor nesting conditions may result in fewer nesting attempts (and thus nests), but does not represent a loss of adults from the breeding population. Instead, the lower number of nests and eggs in a poor nesting year will likely contribute to a reduction in total population size in future years via lower recruitment. This nest survey does not provide precise annual estimates for species that occur in low densities, such as loons, cranes, and swans, or those with clumped distributions, such as brant, common eiders, and gulls. However, long-term averages should be accurate.

A primary advantage of the random nest plot sampling procedure over intensive local studies is that it assures applicability of estimates to the entire coastal zone not just the immediate areas around intensive biological study camps. Moreover, the single brief visit to scattered plots ensures that the monitoring of populations occurs with minimum disturbance. The expansion of estimated nests and eggs from the ground sampled area to the entire coastal zone is based the assumption that observed breeding bird indices obtained from aerial surveys provide an accurate linear relationship with the number of nests within versus outside of the ground sampled area. By using a 7-year localized average Out/In ratio, the variation in aerial estimates due to sampling error is moderated.

Annual changes in nest population size are less informative than long-term trends because of sampling error, changes in observers, distribution of plots, and small sample size for less common species. Only several years of consistent declines or increases are likely to indicate a true change in the number of nests and eggs produced on the Yukon-Kuskokwim coastal zone. We believe that a graphical presentation (Fig. 2) enables better interpretation of data than analysis of year-to-year changes in population size. Large annual changes in nest population size probably reflect sampling error or result from extremes in nesting effort and success, rather than real population change.



ACKNOWLEDGMENTS

The nest plot survey is a cooperative project between Region 7 Migratory Bird Management (MBM) and the Yukon Delta National Wildlife Refuge (YDNWR). Special thanks go to Thomas Doolittle (YDNWR) and Doug Staller (Kenai NWR, formerly with YDNWR) for financial support and logistical coordination. We also thank Victor Anvil (YDNWR) for maintenance of essential facilities and equipment used by the nest plot crews. We are indebted to Hollis Twitchell (YDNWR), Robert Sundown (YDNWR), and Chuck Moore (Regional Aviation Management- RAM) for safe and superb flying during the course of the survey; and special thanks go to George Walters (YDNWR) whose aviation skills, project support, and good cheer have contributed significantly to the success of this project for many years. We also thank Bryce Lake (YDNWR), Louie Andrew (YDNWR), Jim Sedinger (University of Nevada, Reno- UNR), Amanda Walker (UNR), Joel Schmutz (USGS), Patrick Lemons (USGS), Margaret Petersen (USGS), and Matt Sexson (USGS) for providing personnel during their already busy field seasons. We also thank Paul Anderson (MBM) for air support of personnel and supplies, and Bill Eldridge (MBM), Karen Bollinger (MBM), and Bob Platte (MBM) for aerial survey data. Bob Platte also provided invaluable GIS data processing, plot delineation, and preparation of photomaps. We thank Bill Larned, Tim Bowman, and Bob Platte for critical reviews of this report. The following individuals collected data in 2008: Stan Afcan, Doug Alcorn, Mike Blom, Chris Dau, Lynn Denlinger, Tasha DiMarzio, Tyrone Donnelly, Julian Fischer, Paul Flint, Barbara Heindl, Bryce Lake, Susan Lawrence, David Leal, Patrick Lemons, Jessica Lewis, Sarah Lovivond, Dennis Marks, Sarah McCloskey, David Messmer, Tim Moser, Russ Oates, Steve Olson, Bill Ostrand, Anthony Pagano, Molly Schreiner, Rob Scuckard, Matt Sexson, Terry Sharpe, Lynn Snoddy, Matt Sorum, Heather Wilson, and Tamara Zeller.



LITERATURE CITED

- Anthony, R. M., P. L. Flint, and J. S. Sedinger. 1991. Arctic fox removal improves nest success of black brant. *Wildl. Soc. Bull.* 19: 176-184.
- Bollinger, K. S. and W. D. Eldridge. 2008. Report to the Pacific Flyway Committee on the 1985-2008 Coastal Zone Yukon-Kuskokwim Delta Goose Survey of geese, swans and sandhill cranes. Memo. USFWS, Fairbanks, AK.
- Bowman, T. D. 2008. Field guide to bird nests and eggs of Alaska's coastal tundra, 2nd Edition. Alaska Sea Grant College Program, University of Alaska, Fairbanks.
- Bowman, T. D. and R. A. Stehn. *Manuscript in prep.* Nest detection rate on plots searched to monitor Yukon-Kuskokwim Delta waterbird populations.
- Butler, W. I. Jr. 1983. U.S. Fish and Wildlife Service Memorandum, July 19, 1983. Cackling Canada goose nesting population. Yukon Delta NWR.
- Butler, W. I., Jr., R. A. Stehn, and W. D. Eldridge. 1988. Development of an aerial breeding pair survey for geese nesting in the coastal zone of the Yukon Delta. Annual progress report, U.S. Fish and Wildlife Service, Anchorage.
- Dau, C. P. and P. G. Mickelson. 1979. Relation of weather to spring migration and nesting of Cackling Geese on the Yukon-Kuskokwim Delta, pgs. 94-104, *In* R. L. Jarvis and J. C. Bartonek [eds], Management and biology of Pacific Flyway geese. Oregon State University Book Stores. Corvallis, Oregon.
- Fischer, J. B., R. A. Stehn, and G. Walters. 2007. Nest populations and potential production of geese and spectacled eiders on the Yukon-Kuskokwim Delta, Alaska, 2007. Unpubl. Rep., U.S. Fish and Wildlife Service, Anchorage, AK.
- Lake, B. C. 2008. Nesting Ecology of spectacled and common eiders on Kigigak Island, Yukon Delta NWR, Alaska, 2008. Unpubl. Rep. Yukon Delta NWR, Bethel, AK.
- NOAA. 2008. National Weather Service, Alaska-Pacific, River Forecast Center Home Page. <<http://aprfc.arh.noaa.gov/data/breakup.html>>.
- Pamplin, W. L. Jr. 1986. Cooperative efforts to halt population declines of geese nesting on Alaska's Yukon-Kuskokwim Delta. *Trans. N. Am. Wildl. Nat. Resour. Conf.* 51:487-506.
- Platte, R. M., and R. A. Stehn. 2008. Abundance and Trend of Waterbirds on Alaska's Yukon-Kuskokwim Delta Coast based on 1988 to 2008 Aerial Surveys. Unpubl. Rep., U.S. Fish and Wildlife Service, Anchorage, AK.
- Raveling, D. G. 1978. Timing of egg laying by northern geese. *Auk* 95: 294-303.
- Stehn, R. A., W. W. Larned, R. M. Platte, J. B. Fischer, and T. D. Bowman. 2006. Spectacled eider population status and trend in Alaska. Unpubl. Rep., U.S. Fish and Wildlife Service, Anchorage, AK.
- Westerkov, K. 1950. Methods for determining the age of game bird eggs. *J. Wildl. Manage.* 14:56-67.
- Wilson, H. M. 2008. Aerial photographic survey of brant colonies on the Yukon-Kuskokwim Delta, Alaska, 2008. Unpubl. Rep., U.S. Fish and Wildlife Service, Anchorage, AK.

Figure 2 (Subsequent pages). Population size (\pm 90% CI) and trends of waterbird nests and egg production on the Yukon-Kuskokwim Delta Alaska, 1985-2008, with accompanying tabulated data. Column heading definitions follow:

Year = survey year;

N plots = number of ground sampled plots used in the analysis;

Sampled km² = total area searched (N plots*plot size);

Nest index IN = number of nests within the core 716 km² ground sampled area uncorrected for nest detection;

SE nest index IN = standard error for nest index;

Avg nest detection rates = annual proportion of nests detected based on predictive model (considers species, nest status, habitat, observer experience, Bowman and Stehn *manuscript in prep*);

Corrected nest IN = Nest index corrected for nest detection;

7 yr avg aerial Out:In ratio = the seven-year localized average ratio of aerial observations seen out of the ground sampled area vs. in the ground sampled area (seven years are based on from the current year, 3 prior years, and 3 following years data as available);

Corrected nests OUT = number of nests extrapolated beyond the ground sampled area based on the 7-yr localized average Out:In ratio, corrected for nest detection rate;

Total nests In+Out = total number of nests in the YKD coastal zone, corrected for nest detection rate;

SE total nests = standard error for total nest estimate;

Total eggs In+Out = total number of viable eggs at time of plot search in the YKD coastal zone, corrected for detection rate;

SE total eggs = standard error for total egg estimate;


Total eggs/active nests = total viable eggs In+Out divided by the nests with eggs In+Out, corrected for detection rate;

Total eggs/total nests = total viable eggs In+Out divided by the total nests In+Out, corrected for detection rate;


Corrected % nest success = number of active nests divided by total nests times 100%, corrected for detection rate;

Relative production rate = Total eggs/total nests divided by the long-term average of total eggs/total nests. This measure describes how the production of eggs per nests compares to the long-term average illustrating the relative contribution of eggs to the population on a per breeding pair basis.

CCGO Cackling Goose




1985-2008 avg annual growth rate= 1.055 (90%c.i.= 1.042-1.069)
 1999-2008 last 10 yrs annual growth rate= 0.998 (90%c.i.= 0.968-1.028)




1985-2008 avg annual growth rate= 1.060 (90%c.i.= 1.038-1.081)
 1999-2008 last 10 yrs annual growth rate= 1.014 (90%c.i.= 0.945-1.083)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	12,788	1,625	79.2%	16,149	1.58	25,513	41,662	4,522	68,715	10,132	3.91	1.65	42%	47%
1986	46	22.16	10,594	1,406	85.0%	12,467	1.52	18,983	31,451	3,349	93,991	12,032	4.89	2.99	61%	85%
1987	37	12.67	14,909	2,693	86.1%	17,319	1.48	25,626	42,945	6,158	206,345	31,217	5.12	4.80	94%	137%
1988	31	10.04	9,342	1,957	83.2%	11,224	1.49	16,725	27,950	4,468	79,592	13,070	4.52	2.85	63%	81%
1989	23	7.45	16,053	3,536	87.3%	18,390	1.37	25,244	43,634	6,883	162,046	28,156	4.85	3.71	77%	106%
1990	33	10.70	18,465	3,890	83.9%	21,997	1.32	29,040	51,037	7,928	171,285	25,876	4.55	3.36	74%	95%
1991	36	11.66	22,840	3,763	86.5%	26,414	1.26	33,387	59,801	7,483	227,526	27,305	4.64	3.80	82%	108%
1992	42	13.39	25,662	4,554	85.3%	30,098	1.22	36,583	66,680	8,983	293,956	41,285	4.82	4.41	91%	125%
1993	47	15.23	22,469	2,877	85.4%	26,323	1.17	30,753	57,076	5,622	221,063	21,490	4.51	3.87	86%	110%
1994	41	13.27	27,391	3,099	85.5%	32,051	1.14	36,555	68,606	6,025	252,233	23,308	4.58	3.68	80%	105%
1995	50	22.56	43,839	5,413	85.9%	51,015	1.07	54,795	105,810	9,303	399,910	35,053	4.46	3.78	85%	107%
1996	54	19.44	39,761	4,827	85.3%	46,617	1.08	50,546	97,162	8,674	389,565	36,565	4.49	4.01	89%	114%
1997	72	23.31	37,516	4,527	86.1%	43,550	1.11	48,431	91,982	7,947	292,993	28,269	4.03	3.19	79%	91%
1998	64	20.71	54,802	6,330	85.1%	64,403	1.14	73,549	137,952	12,199	570,263	52,640	4.47	4.13	92%	118%
1999	53	16.97	46,698	5,561	84.1%	55,508	1.20	66,830	122,339	11,093	404,090	36,943	3.89	3.30	85%	94%
2000	80	25.86	46,279	3,884	87.0%	53,165	1.24	66,059	119,224	7,594	505,617	32,676	4.50	4.24	94%	121%
2001	81	26.23	32,937	3,999	80.7%	40,799	1.28	52,358	93,157	8,606	187,188	19,151	3.64	2.01	55%	57%
2002	84	27.15	40,438	3,989	84.3%	47,948	1.29	61,973	109,921	8,121	402,247	30,175	4.42	3.66	83%	104%
2003	83	26.87	27,323	2,905	82.6%	33,071	1.31	43,232	76,303	5,831	183,773	17,127	3.96	2.41	61%	69%
2004	81	26.22	36,574	3,024	87.5%	41,818	1.31	54,683	96,501	5,939	387,103	26,534	4.72	4.01	85%	114%
2005	83	26.87	35,666	3,192	87.2%	40,898	1.32	53,920	94,818	6,283	344,377	24,329	4.27	3.63	85%	103%
2006	75	24.28	35,842	3,708	85.9%	41,706	1.29	53,781	95,486	7,331	347,816	28,637	4.43	3.64	82%	104%
2007	79	25.58	46,112	4,684	86.2%	53,492	1.30	69,581	123,073	9,908	519,646	43,879	4.60	4.22	92%	120%
2008	82	26.55	42,566	3,963	83.7%	50,846	1.29	65,439	116,285	8,153	351,091	26,272	4.06	3.02	74%	86%

EMGO Emperor Goose




1985-2008 avg annual growth rate= 1.007 (90%c.i.= 0.993-1.021)
 1999-2008 last 10 yrs annual growth rate= 1.025 (90%c.i.= 0.977-1.072)




1985-2008 avg annual growth rate= 1.010 (90%c.i.= 0.992-1.029)
 1999-2008 last 10 yrs annual growth rate= 1.037 (90%c.i.= 0.973-1.101)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	Tyr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	3,816	685	86.5%	4,411	3.09	13,638	18,049	2,591	60,298	12,169	5.67	3.34	59%	76%
1986	46	22.16	5,426	620	89.0%	6,096	3.01	18,333	24,428	2,317	87,871	8,967	4.93	3.60	73%	81%
1987	37	12.67	8,979	1,477	87.9%	10,218	2.97	30,359	40,577	5,440	199,734	29,156	5.12	4.92	96%	111%
1988	31	10.04	4,920	965	89.0%	5,530	2.87	15,872	21,402	3,386	88,112	13,806	4.63	4.12	89%	93%
1989	23	7.45	11,824	1,769	88.9%	13,306	2.76	36,738	50,044	6,194	235,985	30,641	5.12	4.72	92%	107%
1990	33	10.70	10,704	1,299	85.7%	12,490	2.62	32,779	45,269	5,000	189,853	21,316	4.91	4.19	85%	95%
1991	36	11.66	12,157	1,812	88.4%	13,758	2.45	33,711	47,469	6,131	224,513	29,936	4.89	4.73	97%	107%
1992	42	13.39	10,265	1,372	86.2%	11,906	2.29	27,216	39,122	4,301	191,003	22,297	5.07	4.88	96%	110%
1993	47	15.23	9,777	1,116	84.5%	11,571	2.18	25,203	36,775	3,490	165,442	16,560	4.78	4.50	94%	102%
1994	41	13.27	13,372	1,647	85.9%	15,561	2.08	32,403	47,964	4,544	224,440	23,628	4.99	4.68	94%	106%
1995	50	22.56	9,738	1,127	85.5%	11,389	2.03	23,082	34,471	3,069	161,656	14,375	4.86	4.69	96%	106%
1996	54	19.44	11,008	1,105	85.6%	12,866	2.07	26,635	39,501	3,182	190,198	16,670	5.14	4.81	94%	109%
1997	72	23.31	7,368	736	87.1%	8,461	2.16	18,282	26,743	2,219	121,774	10,551	4.78	4.55	95%	103%
1998	64	20.71	9,295	964	86.7%	10,719	2.20	23,595	34,314	2,875	151,635	13,475	4.64	4.42	95%	100%
1999	53	16.97	10,166	875	86.2%	11,794	2.33	27,515	39,309	3,319	160,030	14,259	4.44	4.07	92%	92%
2000	80	25.86	9,715	929	86.9%	11,185	2.41	26,908	38,093	3,335	186,005	16,620	4.98	4.88	98%	110%
2001	81	26.23	4,503	478	86.4%	5,209	2.44	12,694	17,903	1,651	64,939	7,457	4.81	3.63	75%	82%
2002	84	27.15	8,699	942	85.8%	10,142	2.52	25,532	35,674	3,256	164,016	15,009	4.98	4.60	92%	104%
2003	83	26.87	7,057	768	84.9%	8,311	2.52	20,950	29,261	2,644	109,668	11,862	4.79	3.75	78%	85%
2004	81	26.22	9,690	909	87.7%	11,051	2.45	27,127	38,178	3,065	176,789	14,696	4.88	4.63	95%	105%
2005	83	26.87	10,948	812	87.0%	12,588	2.48	31,160	43,748	2,945	203,005	13,514	4.97	4.64	93%	105%
2006	75	24.28	9,373	957	88.0%	10,648	2.36	25,169	35,817	2,833	162,856	12,345	4.78	4.55	95%	103%
2007	79	25.58	10,241	976	87.6%	11,688	2.35	27,484	39,171	3,001	188,813	15,939	4.98	4.82	97%	109%
2008	82	26.55	9,570	782	86.2%	11,103	2.36	26,154	37,257	2,458	163,478	10,383	4.81	4.39	91%	99%

WFGO White-fronted Goose




1985-2008 avg annual growth rate= 1.097 (90%c.i.= 1.080-1.114)
 1999-2008 last 10 yrs annual growth rate= 1.048 (90%c.i.= 1.023-1.074)




1985-2008 avg annual growth rate= 1.097 (90%c.i.= 1.077-1.118)
 1999-2008 last 10 yrs annual growth rate= 1.049 (90%c.i.= 1.014-1.084)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	1,078	257	81.0%	1,331	5.85	7,789	9,120	2,199	26,944	7,565	4.20	2.95	70%	71%
1986	46	22.16	2,907	463	83.2%	3,493	5.92	20,686	24,179	4,031	94,114	16,817	4.37	3.89	89%	94%
1987	37	12.67	3,275	629	81.4%	4,026	5.88	23,681	27,707	5,107	130,420	23,580	4.90	4.71	96%	114%
1988	31	10.04	2,995	599	83.9%	3,568	6.02	21,488	25,056	4,636	105,590	20,103	4.32	4.21	97%	102%
1989	23	7.45	4,037	1,004	84.9%	4,753	6.00	28,516	33,269	7,624	157,870	34,264	5.01	4.75	95%	115%
1990	33	10.70	7,025	1,108	81.0%	8,674	5.95	51,573	60,247	9,448	248,679	39,565	4.45	4.13	93%	100%
1991	36	11.66	7,184	1,009	86.1%	8,345	5.52	46,042	54,388	6,969	232,734	29,649	4.53	4.28	94%	103%
1992	42	13.39	8,019	1,001	82.6%	9,710	5.49	53,286	62,996	7,469	278,756	34,249	4.52	4.42	98%	107%
1993	47	15.23	5,641	853	80.4%	7,015	5.25	36,816	43,830	6,198	181,874	24,937	4.27	4.15	97%	100%
1994	41	13.27	8,789	1,097	81.3%	10,813	5.12	55,348	66,161	7,956	279,779	34,062	4.34	4.23	97%	102%
1995	50	22.56	9,992	1,093	81.0%	12,340	4.99	61,563	73,903	7,411	303,858	31,087	4.26	4.11	97%	99%
1996	54	19.44	12,849	1,303	82.6%	15,558	5.31	82,587	98,145	9,389	424,347	41,487	4.49	4.32	96%	105%
1997	72	23.31	11,298	1,145	81.7%	13,823	5.49	75,917	89,740	8,785	377,093	39,272	4.28	4.20	98%	102%
1998	64	20.71	12,785	1,320	81.7%	15,657	5.91	92,579	108,236	12,325	456,545	52,664	4.33	4.22	97%	102%
1999	53	16.97	10,588	1,157	82.4%	12,853	6.16	79,213	92,065	10,685	360,806	42,205	4.17	3.92	94%	95%
2000	80	25.86	13,646	1,258	82.9%	16,461	6.37	104,876	121,337	11,868	524,331	52,672	4.45	4.32	97%	104%
2001	81	26.23	11,407	935	82.8%	13,775	6.52	89,869	103,645	8,496	371,884	33,434	3.86	3.59	93%	87%
2002	84	27.15	11,994	1,001	81.6%	14,694	6.33	92,956	107,650	9,961	461,687	44,778	4.39	4.29	98%	104%
2003	83	26.87	11,265	1,151	81.8%	13,773	6.32	87,106	100,878	10,708	402,433	44,274	4.25	3.99	94%	96%
2004	81	26.22	17,059	1,465	82.7%	20,638	6.28	129,507	150,145	14,194	666,798	62,581	4.59	4.44	97%	107%
2005	83	26.87	18,432	1,472	82.2%	22,421	5.88	131,866	154,287	12,342	624,702	51,201	4.24	4.05	96%	98%
2006	75	24.28	17,685	1,571	82.1%	21,537	5.72	123,276	144,812	12,637	573,811	50,287	4.15	3.96	96%	96%
2007	79	25.58	18,579	1,518	84.4%	22,017	5.67	124,842	146,858	12,704	645,241	56,817	4.47	4.39	98%	106%
2008	82	26.55	16,175	1,124	80.8%	20,010	5.63	112,614	132,624	11,707	496,734	45,329	4.05	3.75	92%	91%

BRAN Black Brant




1985-2008 avg annual growth rate= 0.996 (90%c.i.= 0.970-1.023)
 1999-2008 last 10 yrs annual growth rate= 0.983 (90%c.i.= 0.913-1.053)




1985-2008 avg annual growth rate= 1.000 (90%c.i.= 0.961-1.039)
 1999-2008 last 10 yrs annual growth rate= 1.018 (90%c.i.= 0.886-1.150)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	7,107	2,569	92.6%	7,675	3.08	23,626	31,300	9,138	13,080	4,941	3.09	0.42	14%	16%
1986	46	22.16	3,844	2,304	95.7%	4,017	3.08	12,367	16,384	7,812	48,188	29,690	3.99	2.94	74%	115%
1987	37	12.67	13,497	6,158	95.6%	14,117	3.47	49,042	63,159	24,303	264,675	103,411	4.26	4.19	98%	163%
1988	31	10.04	2,852	1,620	95.9%	2,973	3.47	10,330	13,304	6,207	47,365	24,074	3.98	3.56	89%	139%
1989	23	7.45	6,537	2,701	94.8%	6,893	3.45	23,755	30,648	10,338	78,581	30,707	3.49	2.56	73%	100%
1990	33	10.70	8,563	4,710	94.7%	9,047	3.82	34,557	43,604	20,163	80,288	35,026	3.15	1.84	58%	72%
1991	36	11.66	7,859	6,513	94.3%	8,335	3.86	32,201	40,536	27,870	102,937	75,723	3.66	2.54	69%	99%
1992	42	13.39	19,835	9,859	95.6%	20,742	3.97	82,434	103,176	43,232	381,036	163,092	3.87	3.69	95%	144%
1993	47	15.23	14,196	6,832	95.7%	14,838	3.95	58,667	73,505	29,796	172,715	61,309	3.23	2.35	73%	92%
1994	41	13.27	8,681	5,693	96.0%	9,047	3.76	34,004	43,051	23,186	96,524	44,514	2.42	2.24	93%	87%
1995	50	22.56	6,186	3,119	96.5%	6,410	3.56	22,850	29,260	12,241	62,530	21,223	2.98	2.14	72%	83%
1996	54	19.44	4,050	2,022	95.6%	4,235	3.49	14,794	19,029	7,882	61,664	28,011	3.75	3.24	87%	126%
1997	72	23.31	3,807	1,423	96.7%	3,938	3.03	11,947	15,885	4,857	49,158	15,777	3.32	3.09	93%	121%
1998	64	20.71	16,862	5,452	95.3%	17,702	3.08	54,517	72,219	19,316	229,935	65,075	3.67	3.18	87%	124%
1999	53	16.97	6,581	2,064	94.1%	6,991	2.82	19,738	26,729	6,794	68,162	18,018	3.16	2.55	81%	99%
2000	80	25.86	15,140	5,069	96.6%	15,679	2.74	42,891	58,570	15,558	149,228	35,681	2.87	2.55	89%	99%
2001	81	26.23	8,487	2,391	92.7%	9,156	2.45	22,437	31,593	7,060	48,314	13,621	2.95	1.53	52%	60%
2002	84	27.15	11,177	4,344	94.8%	11,792	2.50	29,444	41,235	12,502	112,171	36,456	3.12	2.72	87%	106%
2003	83	26.87	8,229	4,048	94.1%	8,741	2.42	21,146	29,887	11,069	17,457	6,087	1.25	0.58	47%	23%
2004	81	26.22	4,968	1,710	95.7%	5,192	2.36	12,229	17,421	4,682	44,262	12,960	3.28	2.54	77%	99%
2005	83	26.87	10,015	2,732	96.4%	10,385	2.18	22,660	33,045	6,920	83,131	17,624	3.02	2.52	83%	98%
2006	75	24.28	5,541	1,993	95.4%	5,810	2.16	12,522	18,333	5,022	43,432	11,723	3.44	2.37	69%	92%
2007	79	25.58	13,711	4,083	96.5%	14,214	2.05	29,142	43,355	9,870	161,406	37,308	3.94	3.72	95%	145%
2008	82	26.55	11,619	4,337	95.5%	12,169	2.14	26,029	38,198	10,673	94,586	32,615	3.20	2.48	77%	97%

TUSW Tundra Swan




1985-2008 avg annual growth rate= 1.020 (90%c.i.= 1.010-1.030)
 1999-2008 last 10 yrs annual growth rate= 1.013 (90%c.i.= 0.979-1.047)




1985-2008 avg annual growth rate= 1.020 (90%c.i.= 1.006-1.035)
 1999-2008 last 10 yrs annual growth rate= 1.027 (90%c.i.= 0.958-1.096)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	699	119	81.0%	863	5.74	4,953	5,815	889	19,731	3,437	3.93	3.39	86%	90%
1986	46	22.16	1,227	166	80.4%	1,527	5.90	9,008	10,535	1,284	28,989	4,316	3.21	2.75	86%	73%
1987	37	12.67	847	189	82.5%	1,027	6.09	6,260	7,287	1,454	25,481	5,118	3.50	3.50	100%	93%
1988	31	10.04	1,070	247	78.9%	1,355	6.18	8,370	9,725	2,011	44,800	9,512	4.61	4.61	100%	123%
1989	23	7.45	769	263	88.5%	869	6.31	5,481	6,351	1,909	26,383	8,532	4.15	4.15	100%	111%
1990	33	10.70	1,204	288	76.3%	1,579	6.35	10,028	11,607	2,472	42,542	9,358	3.92	3.67	93%	98%
1991	36	11.66	798	178	81.8%	976	6.55	6,398	7,374	1,469	28,668	6,131	4.25	3.89	92%	104%
1992	42	13.39	962	174	83.5%	1,152	6.73	7,759	8,912	1,437	35,204	5,876	3.95	3.95	100%	105%
1993	47	15.23	846	226	79.6%	1,063	6.95	7,384	8,446	2,019	32,587	8,316	4.07	3.86	95%	103%
1994	41	13.27	1,024	231	82.8%	1,237	7.01	8,670	9,907	1,982	41,254	8,093	4.16	4.16	100%	111%
1995	50	22.56	730	135	82.1%	889	7.25	6,446	7,335	1,244	25,261	4,525	3.79	3.44	91%	92%
1996	54	19.44	1,141	177	82.2%	1,389	7.52	10,449	11,837	1,751	55,533	8,002	4.89	4.69	96%	125%
1997	72	23.31	1,074	155	81.0%	1,326	7.95	10,544	11,870	1,607	46,701	6,620	4.20	3.93	94%	105%
1998	64	20.71	1,140	182	81.7%	1,396	8.00	11,175	12,571	1,878	35,119	5,980	3.10	2.79	90%	74%
1999	53	16.97	759	145	82.2%	924	8.25	7,621	8,545	1,488	27,203	5,349	3.63	3.18	88%	85%
2000	80	25.86	913	153	84.2%	1,085	8.23	8,923	10,008	1,541	33,826	5,337	3.62	3.38	93%	90%
2001	81	26.23	819	134	83.1%	986	8.36	8,239	9,225	1,395	25,615	4,083	3.38	2.78	82%	74%
2002	84	27.15	1,054	166	80.9%	1,303	8.21	10,705	12,008	1,747	46,528	7,332	4.35	3.87	89%	103%
2003	83	26.87	1,198	187	82.7%	1,449	8.29	12,002	13,451	1,990	62,625	9,480	4.79	4.66	97%	124%
2004	81	26.22	1,337	189	83.2%	1,608	8.37	13,456	15,064	2,009	80,724	10,641	5.48	5.36	98%	143%
2005	83	26.87	1,039	150	82.5%	1,259	8.70	10,952	12,210	1,657	41,814	5,857	3.63	3.42	94%	91%
2006	75	24.28	884	143	83.3%	1,062	8.76	9,302	10,364	1,579	34,112	5,392	3.54	3.29	93%	88%
2007	79	25.58	895	154	83.6%	1,071	8.94	9,575	10,646	1,728	39,813	6,716	3.87	3.74	97%	100%
2008	82	26.55	782	127	83.3%	939	9.07	8,510	9,449	1,502	33,534	5,846	3.85	3.55	92%	95%

SACR Sandhill Crane




1985-2008 avg annual growth rate= 1.009 (90%c.i.= 0.990-1.028)
 1999-2008 last 10 yrs annual growth rate= 0.996 (90%c.i.= 0.921-1.071)




1985-2008 avg annual growth rate= 1.016 (90%c.i.= 0.995-1.038)
 1999-2008 last 10 yrs annual growth rate= 1.031 (90%c.i.= 0.941-1.122)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	553	145	63.3%	875	9.99	8,740	9,615	2,635	12,605	3,521	1.63	1.31	80%	78%
1986	46	22.16	775	188	74.5%	1,040	9.99	10,392	11,433	2,796	13,150	3,447	1.48	1.15	78%	68%
1987	37	12.67	904	192	62.8%	1,439	9.99	14,374	15,812	3,647	24,958	5,882	1.58	1.58	100%	93%
1988	31	10.04	1,141	225	67.2%	1,699	9.96	16,924	18,623	3,789	37,247	7,578	2.14	2.00	94%	118%
1989	23	7.45	385	178	73.2%	526	9.77	5,135	5,661	2,465	11,322	4,931	2.00	2.00	100%	118%
1990	33	10.70	1,606	305	59.5%	2,696	9.60	25,883	28,580	5,829	48,540	10,616	1.79	1.70	95%	100%
1991	36	11.66	982	222	72.8%	1,350	9.35	12,627	13,978	3,238	26,476	6,376	1.89	1.89	100%	112%
1992	42	13.39	1,283	267	68.2%	1,881	9.48	17,825	19,706	3,959	32,976	6,610	1.75	1.67	96%	99%
1993	47	15.23	893	227	51.8%	1,723	8.83	15,212	16,935	3,930	24,875	5,701	1.67	1.47	88%	87%
1994	41	13.27	1,240	254	62.0%	2,001	8.54	17,096	19,097	3,744	34,262	7,202	1.79	1.79	100%	106%
1995	50	22.56	983	154	49.1%	2,003	8.72	17,467	19,469	3,498	28,855	5,003	1.92	1.48	77%	88%
1996	54	19.44	1,362	213	62.2%	2,191	8.90	19,501	21,692	3,392	37,840	6,087	1.88	1.74	93%	103%
1997	72	23.31	1,044	187	59.8%	1,746	9.50	16,593	18,340	3,161	32,379	5,916	1.77	1.77	100%	104%
1998	64	20.71	1,071	175	58.2%	1,839	10.18	18,725	20,564	3,518	36,414	6,158	1.77	1.77	100%	105%
1999	53	16.97	633	162	44.7%	1,416	10.75	15,222	16,638	4,521	18,562	5,486	1.69	1.12	66%	66%
2000	80	25.86	969	139	62.0%	1,563	10.90	17,039	18,602	2,857	31,621	4,994	1.84	1.70	92%	101%
2001	81	26.23	355	111	65.4%	542	11.01	5,968	6,510	1,884	9,970	3,125	1.70	1.53	90%	91%
2002	84	27.15	1,054	149	54.6%	1,933	10.58	20,453	22,386	3,497	38,666	6,007	2.00	1.73	86%	102%
2003	83	26.87	1,092	155	67.9%	1,608	10.86	17,465	19,073	2,796	36,465	5,531	1.96	1.91	97%	113%
2004	81	26.22	1,256	161	62.7%	2,003	10.62	21,278	23,280	3,211	45,210	6,126	1.94	1.94	100%	115%
2005	83	26.87	1,145	164	58.4%	1,962	10.50	20,601	22,564	3,348	42,619	6,270	1.89	1.89	100%	112%
2006	75	24.28	648	141	61.6%	1,052	10.12	10,648	11,700	2,578	20,602	4,918	1.76	1.76	100%	104%
2007	79	25.58	811	147	66.8%	1,215	10.20	12,391	13,606	2,449	26,781	4,864	1.97	1.97	100%	116%
2008	82	26.55	728	136	60.7%	1,199	10.64	12,760	13,959	2,711	23,541	4,616	1.76	1.69	96%	100%

SPEI Spectacled Eider




1985-2008 avg annual growth rate= 0.987 (90%c.i.= 0.973-1.000)
 1999-2008 last 10 yrs annual growth rate= 1.046 (90%c.i.= 1.001-1.091)




1985-2008 avg annual growth rate= 0.994 (90%c.i.= 0.976-1.013)
 1999-2008 last 10 yrs annual growth rate= 1.043 (90%c.i.= 0.969-1.117)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	2,272	489	77.8%	2,919	1.11	3,238	6,157	966	10,535	2,061	3.84	1.71	45%	44%
1986	46	22.16	2,164	366	77.7%	2,786	1.11	3,090	5,876	747	14,217	2,325	4.40	2.42	55%	63%
1987	37	12.67	3,558	758	77.9%	4,568	1.11	5,067	9,635	1,508	40,284	6,821	5.06	4.18	83%	108%
1988	31	10.04	1,854	500	82.0%	2,261	1.11	2,508	4,770	924	20,859	3,997	4.81	4.37	91%	113%
1989	23	7.45	2,307	751	78.8%	2,927	1.15	3,362	6,289	1,469	28,366	7,148	4.99	4.51	90%	117%
1990	33	10.70	2,141	552	79.6%	2,689	1.01	2,716	5,404	1,071	24,236	5,251	5.03	4.48	89%	116%
1991	36	11.66	1,596	491	76.9%	2,075	0.93	1,940	4,015	938	16,402	4,011	5.27	4.09	77%	106%
1992	42	13.39	1,230	308	77.5%	1,587	0.81	1,289	2,876	565	13,997	2,886	5.43	4.87	90%	126%
1993	47	15.23	1,410	348	75.3%	1,874	0.74	1,385	3,259	628	11,135	2,524	4.27	3.42	80%	89%
1994	41	13.27	1,779	344	77.4%	2,300	0.70	1,607	3,907	638	13,769	2,403	4.65	3.52	76%	91%
1995	50	22.56	2,094	417	78.0%	2,684	0.63	1,679	4,363	690	16,236	2,955	4.88	3.72	76%	97%
1996	54	19.44	1,988	377	77.3%	2,573	0.53	1,373	3,946	571	15,913	2,474	5.06	4.03	80%	105%
1997	72	23.31	1,719	404	82.7%	2,079	0.55	1,154	3,233	546	12,335	2,376	4.38	3.82	87%	99%
1998	64	20.71	2,384	374	80.6%	2,956	0.55	1,623	4,579	537	20,493	2,313	4.87	4.48	92%	116%
1999	53	16.97	2,320	532	81.0%	2,864	0.53	1,513	4,377	749	19,556	3,234	4.96	4.47	90%	116%
2000	80	25.86	1,965	295	82.0%	2,398	0.53	1,260	3,657	422	18,103	2,186	5.34	4.95	93%	128%
2001	81	26.23	1,474	275	78.7%	1,873	0.53	1,002	2,875	402	7,044	1,087	4.18	2.45	59%	64%
2002	84	27.15	2,135	407	80.1%	2,664	0.49	1,300	3,964	568	16,690	3,015	5.26	4.21	80%	109%
2003	83	26.87	1,651	350	82.7%	1,998	0.45	889	2,887	478	9,341	1,883	4.41	3.24	73%	84%
2004	81	26.22	2,102	387	81.1%	2,590	0.43	1,119	3,710	530	14,328	2,325	4.97	3.86	78%	100%
2005	83	26.87	3,489	538	80.3%	4,346	0.43	1,860	6,206	750	25,994	3,645	4.69	4.19	89%	109%
2006	75	24.28	3,272	641	80.6%	4,061	0.45	1,807	5,868	865	20,169	3,187	4.50	3.44	76%	89%
2007	79	25.58	2,490	340	82.7%	3,013	0.43	1,284	4,296	484	18,322	2,276	5.06	4.26	84%	111%
2008	82	26.55	2,911	482	79.5%	3,662	0.36	1,329	4,991	641	19,321	2,821	5.04	3.87	77%	100%

COEI Common Eider




1985-2008 avg annual growth rate= 1.089 (90%c.i.= 1.061-1.117)
 1999-2008 last 10 yrs annual growth rate= 1.109 (90%c.i.= 1.045-1.174)




1985-2008 avg annual growth rate= 1.096 (90%c.i.= 1.061-1.131)
 1999-2008 last 10 yrs annual growth rate= 1.133 (90%c.i.= 1.058-1.208)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	87	47	90.2%	97	1.39	134	231	105	425	322	6.00	1.84	31%	44%
1986	46	22.16	65	45	94.2%	69	1.39	95	164	91	740	416	4.52	4.52	100%	109%
1987	37	12.67	339	152	94.7%	358	1.39	496	854	334	3,822	1,606	5.41	4.47	83%	108%
1988	31	10.04	214	156	86.3%	248	1.39	344	592	341	3,167	1,790	5.35	5.35	100%	129%
1989	23	7.45	1,154	456	91.9%	1,256	1.26	1,580	2,836	959	16,117	5,590	5.68	5.68	100%	137%
1990	33	10.70	335	216	91.2%	367	1.26	462	829	409	3,732	1,841	4.50	4.50	100%	109%
1991	36	11.66	737	381	90.6%	814	1.17	952	1,765	697	4,656	2,118	4.76	2.64	55%	64%
1992	42	13.39	642	254	91.9%	698	0.97	678	1,376	412	6,739	2,159	5.38	4.90	91%	118%
1993	47	15.23	376	203	89.5%	420	1.11	466	886	354	2,359	835	4.43	2.66	60%	64%
1994	41	13.27	431	205	92.7%	465	1.24	578	1,043	361	3,362	1,235	3.73	3.22	86%	78%
1995	50	22.56	539	247	95.1%	567	1.18	671	1,238	411	5,876	2,135	5.08	4.75	93%	115%
1996	54	19.44	773	271	91.4%	846	1.18	996	1,842	487	9,488	2,489	5.44	5.15	95%	125%
1997	72	23.31	737	285	92.1%	800	1.10	878	1,678	477	6,926	2,061	4.53	4.13	91%	100%
1998	64	20.71	1,106	299	87.7%	1,261	1.20	1,513	2,774	568	10,622	2,184	5.01	3.83	76%	93%
1999	53	16.97	717	296	91.7%	782	1.28	1,003	1,785	539	5,987	1,969	4.47	3.35	75%	81%
2000	80	25.86	775	212	92.6%	837	1.16	970	1,807	386	8,501	1,790	4.90	4.70	96%	114%
2001	81	26.23	900	292	93.6%	962	1.02	986	1,947	480	7,160	1,807	4.10	3.68	90%	89%
2002	84	27.15	685	191	92.6%	740	1.07	791	1,531	341	6,933	1,602	4.72	4.53	96%	109%
2003	83	26.87	639	225	92.9%	688	1.20	825	1,513	403	4,736	1,521	4.28	3.13	73%	76%
2004	81	26.22	600	212	94.3%	637	1.28	816	1,453	386	6,635	1,700	4.80	4.57	95%	110%
2005	83	26.87	1,225	298	92.4%	1,325	1.24	1,638	2,964	575	12,621	2,487	4.93	4.26	86%	103%
2006	75	24.28	1,916	751	94.4%	2,030	1.17	2,378	4,408	1,305	18,391	5,608	4.81	4.17	87%	101%
2007	79	25.58	1,763	540	90.5%	1,948	1.30	2,539	4,487	1,073	22,227	5,780	5.22	4.95	95%	120%
2008	82	26.55	1,132	329	91.9%	1,232	1.45	1,792	3,024	670	13,030	3,107	4.82	4.31	89%	104%

RTLO Red-throated Loon




1985-2008 avg annual growth rate= 1.005 (90%c.i.= 0.987-1.023)
 1999-2008 last 10 yrs annual growth rate= 1.024 (90%c.i.= 0.968-1.080)




1985-2008 avg annual growth rate= 1.002 (90%c.i.= 0.984-1.020)
 1999-2008 last 10 yrs annual growth rate= 1.023 (90%c.i.= 0.961-1.085)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	308	70	68.7%	449	8.24	3,697	4,146	1,516	6,158	2,240	1.64	1.49	90%	92%
1986	46	22.16	391	89	74.4%	525	8.24	4,326	4,851	1,750	6,980	2,547	1.64	1.44	88%	89%
1987	37	12.67	530	148	63.9%	829	8.24	6,832	7,661	3,325	13,054	5,170	1.95	1.70	88%	105%
1988	31	10.04	237	69	73.5%	323	8.24	2,661	2,984	1,186	4,362	1,843	1.66	1.46	88%	90%
1989	23	7.45	523	158	72.9%	718	8.24	5,917	6,635	2,619	12,539	4,970	1.89	1.89	100%	117%
1990	33	10.70	481	130	66.6%	722	7.62	5,500	6,223	2,190	10,567	3,814	1.82	1.70	93%	105%
1991	36	11.66	441	106	71.9%	613	7.64	4,686	5,299	1,594	9,153	2,656	1.86	1.73	93%	107%
1992	42	13.39	389	97	66.8%	582	7.45	4,336	4,918	1,421	8,138	2,502	1.86	1.65	89%	102%
1993	47	15.23	358	97	71.8%	498	6.92	3,449	3,947	1,080	7,037	2,041	1.92	1.78	93%	110%
1994	41	13.27	208	78	77.2%	269	7.64	2,055	2,324	826	4,175	1,399	1.80	1.80	100%	111%
1995	50	22.56	378	76	73.0%	518	7.74	4,007	4,524	992	8,171	1,848	1.81	1.81	100%	112%
1996	54	19.44	172	57	71.6%	240	7.37	1,772	2,013	675	3,432	1,201	1.71	1.71	100%	106%
1997	72	23.31	281	67	63.7%	440	7.06	3,110	3,550	955	5,988	1,564	1.91	1.69	88%	104%
1998	64	20.71	526	100	61.5%	855	6.56	5,613	6,468	1,640	9,224	2,193	1.87	1.43	76%	88%
1999	53	16.97	673	131	61.8%	1,089	6.21	6,759	7,847	1,977	11,369	3,099	1.81	1.45	80%	90%
2000	80	25.86	509	89	71.2%	715	5.17	3,698	4,413	999	7,749	1,742	1.89	1.76	93%	109%
2001	81	26.23	476	97	68.4%	695	4.84	3,365	4,060	899	6,055	1,422	1.62	1.49	92%	92%
2002	84	27.15	512	78	71.7%	715	5.53	3,954	4,668	882	7,562	1,503	1.72	1.62	94%	100%
2003	83	26.87	316	70	66.4%	476	5.83	2,773	3,249	824	4,864	1,261	1.73	1.50	86%	93%
2004	81	26.22	355	64	64.9%	547	5.94	3,253	3,801	795	4,992	1,065	1.72	1.31	76%	81%
2005	83	26.87	666	119	74.8%	890	6.05	5,387	6,277	1,266	11,138	2,305	1.81	1.77	98%	110%
2006	75	24.28	477	95	72.0%	663	6.42	4,256	4,919	1,072	7,512	1,702	1.72	1.53	89%	95%
2007	79	25.58	594	109	73.4%	808	6.88	5,566	6,374	1,394	10,246	2,206	1.74	1.61	92%	99%
2008	82	26.55	664	124	69.2%	959	6.57	6,304	7,263	1,793	10,767	2,587	1.64	1.48	91%	92%

PALO Pacific Loon




1985-2008 avg annual growth rate= 1.010 (90%c.i.= 0.993-1.027)
 1999-2008 last 10 yrs annual growth rate= 0.974 (90%c.i.= 0.922-1.027)




1985-2008 avg annual growth rate= 1.007 (90%c.i.= 0.990-1.024)
 1999-2008 last 10 yrs annual growth rate= 0.973 (90%c.i.= 0.915-1.032)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	711	151	68.7%	1,035	21.42	22,159	23,194	6,270	34,450	9,217	1.64	1.49	90%	92%
1986	46	22.16	901	191	74.4%	1,211	21.42	25,930	27,141	7,146	39,048	10,514	1.64	1.44	88%	89%
1987	37	12.67	1,221	326	63.9%	1,912	21.42	40,950	42,862	15,483	73,032	22,784	1.95	1.70	88%	105%
1988	31	10.04	547	153	73.5%	745	21.42	15,948	16,693	5,237	24,402	8,457	1.66	1.46	88%	90%
1989	23	7.45	1,207	351	72.9%	1,656	21.42	35,463	37,119	11,517	70,148	21,918	1.89	1.89	100%	117%
1990	33	10.70	1,125	295	66.6%	1,689	19.37	32,720	34,409	10,427	58,432	18,368	1.82	1.70	93%	105%
1991	36	11.66	1,155	254	71.9%	1,607	18.71	30,067	31,674	8,187	54,708	13,393	1.86	1.73	93%	107%
1992	42	13.39	1,001	236	66.8%	1,498	19.21	28,775	30,273	7,852	50,095	14,122	1.86	1.65	89%	102%
1993	47	15.23	911	235	71.8%	1,269	18.48	23,465	24,734	6,498	44,099	12,431	1.92	1.78	93%	110%
1994	41	13.27	601	219	77.2%	779	19.31	15,041	15,820	5,598	28,424	9,395	1.80	1.80	100%	111%
1995	50	22.56	1,018	179	73.0%	1,395	19.44	27,114	28,508	5,512	51,490	10,408	1.81	1.81	100%	112%
1996	54	19.44	454	141	71.6%	633	22.07	13,977	14,610	5,252	24,912	9,358	1.71	1.71	100%	106%
1997	72	23.31	733	158	63.7%	1,149	22.97	26,407	27,557	7,256	46,476	11,809	1.91	1.69	88%	104%
1998	64	20.71	1,305	225	61.5%	2,122	22.82	48,423	50,545	12,284	72,076	16,105	1.87	1.43	76%	88%
1999	53	16.97	1,689	299	61.8%	2,732	21.23	58,009	60,742	14,940	87,999	23,827	1.81	1.45	80%	90%
2000	80	25.86	1,207	188	71.2%	1,695	19.95	33,814	35,509	8,615	62,344	15,007	1.89	1.76	93%	109%
2001	81	26.23	1,080	211	68.4%	1,578	19.32	30,483	32,061	8,813	47,816	13,806	1.62	1.49	92%	92%
2002	84	27.15	1,201	172	71.7%	1,677	19.79	33,182	34,859	8,060	56,469	13,618	1.72	1.62	94%	100%
2003	83	26.87	749	163	66.4%	1,129	17.53	19,801	20,930	5,050	31,339	7,772	1.73	1.50	86%	93%
2004	81	26.22	764	123	64.9%	1,176	19.46	22,888	24,064	4,916	31,607	6,623	1.72	1.31	76%	81%
2005	83	26.87	1,279	214	74.8%	1,709	20.27	34,658	36,368	7,152	64,533	13,107	1.81	1.77	98%	110%
2006	75	24.28	878	165	72.0%	1,220	21.81	26,606	27,825	5,670	42,492	9,119	1.72	1.53	89%	95%
2007	79	25.58	1,085	179	73.4%	1,478	22.86	33,769	35,247	7,037	56,657	11,066	1.74	1.61	92%	99%
2008	82	26.55	1,089	191	69.2%	1,573	24.21	38,085	39,658	8,401	58,793	11,922	1.64	1.48	91%	92%

GLGU Glaucous Gull




1985-2008 avg annual growth rate= 1.014 (90%c.i.= 0.997-1.030)
 1999-2008 last 10 yrs annual growth rate= 1.031 (90%c.i.= 0.978-1.084)




1985-2008 avg annual growth rate= 1.024 (90%c.i.= 1.006-1.041)
 1999-2008 last 10 yrs annual growth rate= 1.048 (90%c.i.= 1.000-1.096)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	2,330	487	88.8%	2,625	5.34	14,030	16,655	3,219	28,705	6,490	2.08	1.72	83%	76%
1986	46	22.16	1,486	316	89.3%	1,663	5.34	8,891	10,554	2,021	15,984	4,148	2.15	1.51	70%	67%
1987	37	12.67	2,089	766	85.0%	2,457	5.34	13,133	15,590	5,387	30,986	11,016	2.15	1.99	93%	87%
1988	31	10.04	1,498	647	92.2%	1,624	5.34	8,683	10,307	3,879	23,353	9,237	2.53	2.27	90%	100%
1989	23	7.45	2,019	1,106	91.4%	2,208	5.34	11,800	14,008	6,833	27,716	12,208	2.27	1.98	87%	87%
1990	33	10.70	1,606	689	91.3%	1,758	5.34	9,396	11,154	4,102	29,035	11,384	2.91	2.60	90%	114%
1991	36	11.66	2,395	1,501	86.9%	2,754	5.34	14,720	17,475	9,637	33,296	17,837	2.99	1.91	64%	84%
1992	42	13.39	3,582	1,211	88.9%	4,027	5.34	21,524	25,551	7,673	64,122	19,803	2.71	2.51	92%	110%
1993	47	15.23	2,021	703	90.7%	2,228	5.27	11,741	13,969	4,201	34,292	11,007	2.53	2.45	97%	108%
1994	41	13.27	2,319	1,103	91.6%	2,532	5.06	12,816	15,348	6,301	42,132	18,381	2.83	2.75	97%	121%
1995	50	22.56	2,252	643	92.8%	2,428	5.20	12,634	15,062	3,744	38,489	9,426	2.56	2.56	100%	112%
1996	54	19.44	884	241	94.0%	940	4.99	4,689	5,629	1,373	14,029	3,345	2.49	2.49	100%	110%
1997	72	23.31	2,548	1,188	93.8%	2,716	4.76	12,916	15,632	6,235	27,649	7,407	1.79	1.77	99%	78%
1998	64	20.71	3,939	1,749	87.6%	4,495	4.47	20,107	24,601	9,578	63,904	24,556	2.67	2.60	97%	114%
1999	53	16.97	1,603	387	88.9%	1,804	4.48	8,079	9,883	2,143	25,124	5,607	2.72	2.54	94%	112%
2000	80	25.86	3,709	974	91.5%	4,054	4.44	18,005	22,059	5,092	53,612	12,512	2.50	2.43	97%	107%
2001	81	26.23	2,347	955	86.3%	2,718	4.44	12,056	14,774	5,165	32,333	11,743	2.36	2.19	93%	96%
2002	84	27.15	2,531	580	86.7%	2,917	4.06	11,833	14,750	3,025	32,641	6,852	2.59	2.21	85%	97%
2003	83	26.87	3,835	1,748	88.4%	4,338	4.52	19,596	23,934	9,241	31,702	9,774	1.64	1.32	81%	58%
2004	81	26.22	2,320	717	91.5%	2,534	4.48	11,361	13,896	3,886	34,087	9,763	2.49	2.45	99%	108%
2005	83	26.87	3,782	1,049	92.6%	4,084	4.12	16,836	20,920	5,459	49,252	11,938	2.41	2.35	98%	104%
2006	75	24.28	2,446	742	94.1%	2,600	3.94	10,255	12,855	3,715	34,118	10,360	2.69	2.65	99%	117%
2007	79	25.58	4,057	1,101	93.0%	4,360	3.79	16,531	20,891	5,892	58,600	16,788	2.83	2.80	99%	123%
2008	82	26.55	3,558	1,156	91.2%	3,901	3.74	14,607	18,508	6,495	46,603	16,458	2.67	2.52	94%	111%

MEGU Mew Gull




1985-2008 avg annual growth rate= 1.007 (90%c.i.= 0.979-1.034)
 1999-2008 last 10 yrs annual growth rate= 1.067 (90%c.i.= 1.011-1.123)




1985-2008 avg annual growth rate= 1.015 (90%c.i.= 0.987-1.044)
 1999-2008 last 10 yrs annual growth rate= 1.067 (90%c.i.= 1.012-1.123)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	1,107	270	72.6%	1,525	4.14	6,309	7,834	1,873	12,599	3,582	1.95	1.61	82%	69%
1986	46	22.16	937	286	80.9%	1,157	4.14	4,790	5,947	1,772	12,887	3,977	2.29	2.17	95%	93%
1987	37	12.67	847	385	67.8%	1,249	4.14	5,168	6,417	2,534	10,019	4,617	1.56	1.56	100%	67%
1988	31	10.04	2,496	1,871	65.9%	3,789	4.14	15,681	19,470	11,386	40,651	26,743	2.47	2.09	85%	90%
1989	23	7.45	1,346	385	69.7%	1,930	4.14	7,984	9,914	3,002	26,160	7,575	2.64	2.64	100%	113%
1990	33	10.70	1,338	814	63.9%	2,095	4.14	8,667	10,762	5,230	26,791	13,871	2.68	2.49	93%	107%
1991	36	11.66	1,719	1,037	67.1%	2,562	4.14	10,604	13,166	7,277	22,303	12,481	2.37	1.69	72%	73%
1992	42	13.39	1,176	323	72.9%	1,612	4.14	6,672	8,285	2,292	21,061	6,052	2.69	2.54	95%	109%
1993	47	15.23	3,667	2,931	78.7%	4,661	4.23	19,716	24,377	15,547	68,382	44,008	2.81	2.81	100%	120%
1994	41	13.27	1,024	547	70.7%	1,450	3.93	5,694	7,143	3,395	17,436	9,286	2.65	2.44	92%	105%
1995	50	22.56	1,396	403	81.2%	1,719	3.92	6,745	8,465	2,151	21,904	5,625	2.59	2.59	100%	111%
1996	54	19.44	847	241	78.2%	1,083	3.47	3,763	4,846	1,177	12,787	3,331	2.64	2.64	100%	113%
1997	72	23.31	276	85	85.6%	323	3.42	1,104	1,426	372	3,501	955	2.45	2.45	100%	105%
1998	64	20.71	1,348	446	80.0%	1,685	3.41	5,738	7,423	2,071	18,009	5,040	2.43	2.43	100%	104%
1999	53	16.97	1,476	399	70.7%	2,089	3.26	6,813	8,903	1,970	21,335	5,018	2.48	2.40	97%	103%
2000	80	25.86	1,024	189	72.4%	1,414	3.12	4,411	5,825	1,043	13,966	2,656	2.59	2.40	93%	103%
2001	81	26.23	982	300	61.8%	1,588	3.27	5,201	6,790	1,854	16,523	4,683	2.43	2.43	100%	104%
2002	84	27.15	1,687	378	73.2%	2,305	3.44	7,924	10,229	2,086	26,211	5,332	2.68	2.56	96%	110%
2003	83	26.87	1,465	387	57.8%	2,535	3.72	9,443	11,978	3,048	24,543	5,785	2.58	2.05	79%	88%
2004	81	26.22	1,419	326	73.4%	1,934	3.53	6,829	8,763	1,968	18,640	3,891	2.45	2.13	87%	91%
2005	83	26.87	3,090	1,366	81.0%	3,813	3.59	13,698	17,511	6,486	43,559	17,470	2.49	2.49	100%	107%
2006	75	24.28	2,004	507	76.4%	2,623	3.75	9,837	12,461	2,711	30,801	7,003	2.47	2.47	100%	106%
2007	79	25.58	1,287	252	75.5%	1,705	3.78	6,441	8,147	1,633	21,463	4,348	2.63	2.63	100%	113%
2008	82	26.55	1,968	520	68.7%	2,866	3.85	11,023	13,889	3,775	30,975	8,152	2.49	2.23	90%	96%

SAGU Sabine's Gull




1985-2008 avg annual growth rate= 1.082 (90%c.i.= 1.054-1.111)
 1999-2008 last 10 yrs annual growth rate= 0.979 (90%c.i.= 0.921-1.038)




1985-2008 avg annual growth rate= 1.067 (90%c.i.= 1.041-1.094)
 1999-2008 last 10 yrs annual growth rate= 0.976 (90%c.i.= 0.905-1.047)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	200	101	39.9%	501	2.14	1,069	1,570	685	4,518	1,982	2.88	2.88	100%	123%
1986	46	22.16	129	78	31.5%	410	2.14	875	1,286	666	3,856	1,998	3.00	3.00	100%	128%
1987	37	12.67	113	76	25.9%	436	2.14	931	1,366	706	4,099	2,119	3.00	3.00	100%	128%
1988	31	10.04	357	147	54.3%	656	2.14	1,401	2,057	686	5,600	1,834	2.72	2.72	100%	116%
1989	23	7.45	673	291	32.8%	2,053	2.14	4,384	6,438	2,284	18,442	6,831	2.86	2.86	100%	122%
1990	33	10.70	669	223	33.3%	2,007	2.14	4,285	6,292	1,793	15,517	4,273	2.47	2.47	100%	105%
1991	36	11.66	675	192	31.6%	2,134	2.14	4,557	6,692	1,714	11,139	3,070	1.66	1.66	100%	71%
1992	42	13.39	802	291	28.3%	2,833	2.14	6,049	8,882	2,562	22,779	6,480	2.56	2.56	100%	110%
1993	47	15.23	1,410	724	33.4%	4,225	2.28	9,611	13,836	5,465	30,198	10,716	2.41	2.18	91%	93%
1994	41	13.27	647	220	29.7%	2,179	2.57	5,600	7,779	2,422	20,007	6,557	2.57	2.57	100%	110%
1995	50	22.56	698	185	34.5%	2,024	2.63	5,332	7,356	1,584	18,669	3,973	2.54	2.54	100%	109%
1996	54	19.44	736	216	28.4%	2,591	2.60	6,730	9,321	2,487	17,755	4,343	2.42	1.90	79%	81%
1997	72	23.31	460	136	38.5%	1,196	2.68	3,209	4,405	1,205	9,625	2,903	2.18	2.18	100%	93%
1998	64	20.71	1,486	720	49.1%	3,026	2.92	8,831	11,857	3,799	24,397	7,403	2.06	2.06	100%	88%
1999	53	16.97	1,181	560	37.9%	3,113	3.15	9,809	12,921	3,975	28,388	8,854	2.20	2.20	100%	94%
2000	80	25.86	775	182	32.2%	2,408	3.03	7,295	9,703	2,231	24,834	6,068	2.56	2.56	100%	109%
2001	81	26.23	1,201	423	24.4%	4,915	2.72	13,378	18,293	6,440	37,355	15,425	2.30	2.04	89%	87%
2002	84	27.15	1,239	404	32.8%	3,774	2.55	9,629	13,402	3,190	32,140	7,612	2.40	2.40	100%	103%
2003	83	26.87	692	186	26.1%	2,656	2.60	6,907	9,563	2,297	17,726	3,958	2.18	1.85	85%	79%
2004	81	26.22	600	148	27.3%	2,199	2.45	5,395	7,594	1,708	11,787	2,756	1.99	1.55	78%	66%
2005	83	26.87	1,145	256	32.0%	3,579	2.24	8,013	11,592	2,333	24,651	5,350	2.13	2.13	100%	91%
2006	75	24.28	1,061	372	40.6%	2,616	2.09	5,474	8,091	2,017	18,157	4,867	2.24	2.24	100%	96%
2007	79	25.58	1,819	398	30.4%	5,992	2.11	12,671	18,663	3,825	42,822	8,427	2.29	2.29	100%	98%
2008	82	26.55	944	151	33.5%	2,819	2.17	6,116	8,935	1,349	20,249	3,154	2.27	2.27	100%	97%

ARTE Arctic Tern



1985-2008 avg annual growth rate= 1.080 (90%c.i.= 1.041-1.118)
 1999-2008 last 10 yrs annual growth rate= 1.017 (90%c.i.= 0.906-1.127)



1985-2008 avg annual growth rate= 1.082 (90%c.i.= 1.040-1.124)
 1999-2008 last 10 yrs annual growth rate= 1.013 (90%c.i.= 0.903-1.124)

Year	N plots	Sampled km2	Nest index IN	SE nest index IN	Avg nest detection rate	Corrected nests IN	7yr avg Aerial OUT:IN ratio	Corrected nests OUT	Total nests IN+OUT	SE total nests	Total eggs IN+OUT	SE total eggs	Total eggs / active nests	Total eggs / total nests	Corrected % nest success	Relative production rate
1985	49	24.57	291	110	60.3%	483	9.42	4,545	5,028	1,798	10,056	3,860	2.00	2.00	100%	108%
1986	46	22.16	194	77	66.6%	291	9.42	2,742	3,033	1,131	5,663	2,298	1.87	1.87	100%	101%
1987	37	12.67	113	112	70.1%	161	9.42	1,517	1,678	1,529	1,678	1,529	1.00	1.00	100%	54%
1988	31	10.04	71	71	57.9%	123	9.42	1,159	1,282	1,170	2,565	2,340	2.00	2.00	100%	108%
1989	23	7.45	577	284	54.7%	1,054	9.42	9,922	10,976	4,983	20,448	9,353	1.86	1.86	100%	101%
1990	33	10.70	335	168	60.3%	554	9.42	5,220	5,774	2,712	9,239	4,841	1.60	1.60	100%	86%
1991	36	11.66	123	85	63.3%	194	9.42	1,827	2,021	1,291	2,981	1,984	1.48	1.48	100%	80%
1992	42	13.39	748	235	61.1%	1,225	9.42	11,534	12,758	3,918	24,279	7,164	1.90	1.90	100%	103%
1993	47	15.23	893	482	43.2%	2,066	9.49	19,594	21,660	10,351	49,285	22,617	2.28	2.28	100%	123%
1994	41	13.27	323	163	59.5%	544	9.64	5,239	5,782	2,684	13,466	6,569	2.33	2.33	100%	126%
1995	50	22.56	539	165	56.1%	961	9.41	9,036	9,996	3,030	19,991	6,343	2.00	2.00	100%	108%
1996	54	19.44	221	85	52.5%	421	8.67	3,648	4,068	1,508	7,547	2,892	1.86	1.86	100%	100%
1997	72	23.31	154	78	64.6%	238	8.91	2,118	2,355	1,131	4,710	2,261	2.00	2.00	100%	108%
1998	64	20.71	691	193	51.3%	1,348	8.72	11,747	13,094	3,353	28,923	7,620	2.21	2.21	100%	119%
1999	53	16.97	591	170	49.6%	1,190	8.73	10,390	11,580	3,266	23,034	6,548	1.99	1.99	100%	107%
2000	80	25.86	277	90	58.6%	473	8.52	4,028	4,500	1,431	8,198	2,591	1.82	1.82	100%	98%
2001	81	26.23	682	269	26.0%	2,623	8.06	21,149	23,772	13,624	35,683	17,293	1.92	1.50	78%	81%
2002	84	27.15	1,529	434	37.4%	4,091	8.01	32,781	36,872	10,505	75,148	21,061	2.04	2.04	100%	110%
2003	83	26.87	506	136	49.2%	1,028	8.18	8,409	9,436	2,351	19,180	5,604	2.03	2.03	100%	110%
2004	81	26.22	737	137	41.2%	1,789	7.61	13,609	15,398	3,441	22,248	4,233	1.77	1.44	82%	78%
2005	83	26.87	1,199	258	60.4%	1,985	7.32	14,523	16,509	3,035	27,956	4,635	1.69	1.69	100%	91%
2006	75	24.28	884	175	42.4%	2,087	7.15	14,914	17,000	3,535	32,812	6,700	1.93	1.93	100%	104%
2007	79	25.58	923	169	51.1%	1,806	6.92	12,487	14,293	2,492	26,406	4,970	1.85	1.85	100%	100%
2008	82	26.55	647	170	53.4%	1,213	6.85	8,305	9,518	2,573	16,883	4,519	1.77	1.77	100%	96%




Figure 3. Estimated average hatch date based on egg float angles, 1982-2008. Linear regression on cackling goose hatch date indicates an average advance of 0.4 days per year.




Figure 4. Comparison of spectacled eider apparent nest success measures at Kigigak Island (successful hatched nests/total nests; Lake 2008) and the Yukon-Kuskokwim Delta nest plot survey (active nests at time of search/total nests, corrected for nest detection rate), 1992-2008.




Figure 5. Numbers of black brant nests as measured by an aerial photographic survey in 5 primary colonies (Wilson 2008) and as measured by this survey in the ground sampled area (nest estimates in ground sampled area only, not expanded to entire coastal zone). The aerial photographic survey and ground survey both include the colony at Kigigak Island.

Table 1. Estimated hatch date based on egg float angles. Means calculated using nest as sample unit. Years with fewer than 3 nests per species not included in calculations.

Year	Mean	Min	Max	N	Year	Average	Min	Max	N
Cackling Goose					Emperor Goose				
1982	4-Jul	25-Jun	18-Jul	170	1982	3-Jul	16-Jun	11-Jul	71
1983	25-Jun	15-Jun	14-Jul	284	1983	22-Jun	14-Jun	6-Jul	100
1984	25-Jun	16-Jun	11-Jul	92	1984	23-Jun	16-Jun	2-Jul	43
1985	3-Jul	24-Jun	15-Jul	278	1985	1-Jul	23-Jun	11-Jul	107
1986	29-Jun	13-Jun	15-Jul	346	1986	27-Jun	18-Jun	9-Jul	196
1987	28-Jun	20-Jun	18-Jul	204	1987	27-Jun	18-Jun	7-Jul	141
1988	24-Jun	15-Jun	8-Jul	66	1988	21-Jun	16-Jun	4-Jul	67
1989	1-Jul	22-Jun	10-Jul	55	1989	30-Jun	18-Jun	7-Jul	63
1990	24-Jun	13-Jun	6-Jul	194	1990	21-Jun	11-Jun	6-Jul	99
1991	22-Jun	12-Jun	3-Jul	352	1991	20-Jun	10-Jun	2-Jul	256
1992	30-Jun	20-Jun	21-Jul	391	1992	29-Jun	21-Jun	9-Jul	182
1993	24-Jun	9-Jun	6-Jul	358	1993	21-Jun	11-Jun	4-Jul	139
1994	19-Jun	8-Jun	9-Jul	409	1994	19-Jun	12-Jun	30-Jun	192
1995	20-Jun	11-Jun	5-Jul	725	1995	18-Jun	10-Jun	6-Jul	188
1996	17-Jun	7-Jun	5-Jul	755	1996	16-Jun	4-Jun	23-Jun	185
1997	17-Jun	3-Jun	4-Jul	812	1997	15-Jun	6-Jun	30-Jun	153
1998	25-Jun	12-Jun	9-Jul	889	1998	24-Jun	16-Jun	3-Jul	215
1999	27-Jun	17-Jun	16-Jul	772	1999	27-Jun	17-Jun	6-Jul	188
2000	24-Jun	14-Jun	10-Jul	1014	2000	23-Jun	13-Jun	8-Jul	280
2001	28-Jun	15-Jun	9-Jul	522	2001	27-Jun	19-Jun	2-Jul	104
2002	20-Jun	10-Jun	4-Jul	930	2002	18-Jun	9-Jun	29-Jun	249
2003	17-Jun	3-Jun	4-Jul	562	2003	15-Jun	5-Jun	26-Jun	153
2004	13-Jun	4-Jun	1-Jul	964	2004	12-Jun	4-Jun	24-Jun	253
2005	20-Jun	9-Jun	7-Jul	957	2005	17-Jun	7-Jun	29-Jun	303
2006	27-Jun	15-Jun	8-Jul	849	2006	24-Jun	16-Jun	4-Jul	253
2007	19-Jun	9-Jun	5-Jul	1027	2007	19-Jun	7-Jun	28-Jun	275
2008	22-Jun	11-Jun	7-Jul	903	2008	21-Jun	12-Jun	7-Jul	239
Mean	24-Jun	13-Jun	9-Jul	551	Mean	22-Jun	13-Jun	3-Jul	174
White-fronted Goose					Black Brant				
1982	4-Jul	26-Jun	12-Jul	14	1982	--	--	--	--
1983	21-Jun	13-Jun	19-Jul	25	1983	23-Jun	15-Jun	3-Jul	11
1984	23-Jun	16-Jun	1-Jul	25	1984	20-Jun	19-Jun	20-Jun	4
1985	30-Jun	23-Jun	7-Jul	42	1985	27-Jun	23-Jun	8-Jul	29
1986	26-Jun	17-Jun	12-Jul	102	1986	25-Jun	19-Jun	6-Jul	126
1987	26-Jun	19-Jun	3-Jul	60	1987	25-Jun	22-Jun	3-Jul	167
1988	22-Jun	15-Jun	3-Jul	32	1988	20-Jun	14-Jun	3-Jul	38
1989	28-Jun	22-Jun	4-Jul	21	1989	27-Jun	19-Jun	6-Jul	40
1990	22-Jun	11-Jun	29-Jun	52	1990	21-Jun	15-Jun	1-Jul	119
1991	21-Jun	12-Jun	3-Jul	138	1991	19-Jun	12-Jun	1-Jul	183
1992	29-Jun	19-Jun	24-Jul	110	1992	26-Jun	19-Jun	6-Jul	152
1993	23-Jun	17-Jun	5-Jul	84	1993	21-Jun	12-Jun	27-Jun	107
1994	19-Jun	11-Jun	28-Jun	129	1994	17-Jun	10-Jun	27-Jun	93
1995	21-Jun	9-Jun	1-Jul	178	1995	18-Jun	12-Jun	1-Jul	41
1996	18-Jun	7-Jun	30-Jun	144	1996	17-Jun	11-Jun	26-Jun	44
1997	18-Jun	7-Jun	29-Jun	184	1997	14-Jun	3-Jun	24-Jun	100
1998	25-Jun	17-Jun	6-Jul	261	1998	23-Jun	16-Jun	4-Jul	260
1999	27-Jun	19-Jun	10-Jul	208	1999	26-Jun	17-Jun	7-Jul	108
2000	25-Jun	14-Jun	9-Jul	334	2000	23-Jun	16-Jun	3-Jul	216
2001	28-Jun	19-Jun	7-Jul	311	2001	26-Jun	19-Jun	5-Jul	77
2002	22-Jun	14-Jun	30-Jun	306	2002	19-Jun	6-Jun	3-Jul	163
2003	19-Jun	6-Jun	1-Jul	272	2003	16-Jun	7-Jun	26-Jun	56
2004	15-Jun	4-Jun	27-Jun	364	2004	11-Jun	4-Jun	24-Jun	101
2005	20-Jun	12-Jun	1-Jul	438	2005	17-Jun	6-Jun	26-Jun	148
2006	26-Jun	16-Jun	10-Jul	370	2006	24-Jun	16-Jun	9-Jul	123
2007	19-Jun	8-Jun	2-Jul	446	2007	18-Jun	9-Jun	29-Jun	147
2008	22-Jun	12-Jun	7-Jul	335	2008	20-Jun	13-Jun	30-Jun	103
Mean	23-Jun	14-Jun	5-Jul	185	Mean	21-Jun	14-Jun	1-Jul	106

Table 1. Estimated hatch date continued.

Year	Average	Min	Max	N
Sandhill Crane				
1982	24-Jun	22-Jun	25-Jun	4
1983	26-Jun	17-Jun	11-Jul	14
1984	19-Jun	15-Jun	21-Jun	6
1985	30-Jun	19-Jun	4-Jul	13
1986	27-Jun	16-Jun	9-Jul	25
1987	25-Jun	18-Jun	10-Jul	16
1988	19-Jun	17-Jun	25-Jun	6
1989	--	--	--	--
1990	18-Jun	15-Jun	22-Jun	9
1991	16-Jun	10-Jun	26-Jun	25
1992	30-Jun	24-Jun	5-Jul	9
1993	19-Jun	15-Jun	27-Jun	14
1994	14-Jun	11-Jun	16-Jun	5
1995	18-Jun	12-Jun	30-Jun	10
1996	14-Jun	10-Jun	25-Jun	14
1997	15-Jun	11-Jun	24-Jun	8
1998	21-Jun	15-Jun	26-Jun	19
1999	23-Jun	19-Jun	28-Jun	12
2000	19-Jun	13-Jun	29-Jun	22
2001	21-Jun	19-Jun	23-Jun	7
2002	19-Jun	8-Jun	3-Jul	12
2003	14-Jun	7-Jun	25-Jun	13
2004	15-Jun	9-Jun	22-Jun	10
2005	15-Jun	10-Jun	26-Jun	23
2006	23-Jun	17-Jun	8-Jul	19
2007	12-Jun	7-Jun	24-Jun	16
2008	20-Jun	15-Jun	24-Jun	12
Mean	20-Jun	14-Jun	28-Jun	13

Year	Average	Min	Max	N
Tundra Swan				
1982	5-Jul	23-Jun	14-Jul	11
1983	24-Jun	15-Jun	30-Jun	6
1984	27-Jun	20-Jun	5-Jul	6
1985	4-Jul	26-Jun	10-Jul	14
1986	28-Jun	19-Jun	10-Jul	23
1987	30-Jun	23-Jun	6-Jul	12
1988	27-Jun	17-Jun	4-Jul	4
1989	1-Jul	29-Jun	3-Jul	4
1990	25-Jun	21-Jun	27-Jun	4
1991	24-Jun	17-Jun	8-Jul	12
1992	30-Jun	24-Jun	7-Jul	8
1993	26-Jun	19-Jun	1-Jul	6
1994	22-Jun	13-Jun	30-Jun	9
1995	25-Jun	21-Jun	2-Jul	9
1996	19-Jun	10-Jun	28-Jun	9
1997	21-Jun	14-Jun	25-Jun	13
1998	30-Jun	23-Jun	12-Jul	20
1999	1-Jul	24-Jun	9-Jul	14
2000	26-Jun	18-Jun	5-Jul	22
2001	30-Jun	19-Jun	9-Jul	16
2002	26-Jun	20-Jun	1-Jul	10
2003	18-Jun	11-Jun	24-Jun	21
2004	19-Jun	10-Jun	27-Jun	16
2005	23-Jun	16-Jun	29-Jun	18
2006	1-Jul	22-Jun	8-Jul	14
2007	24-Jun	16-Jun	2-Jul	19
2008	27-Jun	20-Jun	3-Jul	18
Mean	26-Jun	19-Jun	4-Jul	13

Year	Average	Min	Max	N
Spectacled Eider				
1982	8-Jul	30-Jun	22-Jul	18
1983	29-Jun	20-Jun	6-Jul	22
1984	2-Jul	25-Jun	5-Jul	3
1985	4-Jul	26-Jun	18-Jul	20
1986	2-Jul	22-Jun	20-Jul	38
1987	28-Jun	17-Jun	9-Jul	27
1988	26-Jun	20-Jun	2-Jul	19
1989	2-Jul	22-Jun	7-Jul	5
1990	23-Jun	18-Jun	27-Jun	15
1991	22-Jun	16-Jun	10-Jul	25
1992	2-Jul	26-Jun	14-Jul	17
1993	25-Jun	17-Jun	9-Jul	18
1994	23-Jun	12-Jun	6-Jul	15
1995	24-Jun	14-Jun	4-Jul	44
1996	18-Jun	12-Jun	2-Jul	33
1997	19-Jun	11-Jun	30-Jun	39
1998	28-Jun	17-Jun	7-Jul	52
1999	28-Jun	18-Jun	9-Jul	51
2000	28-Jun	18-Jun	9-Jul	52
2001	3-Jul	25-Jun	16-Jul	32
2002	22-Jun	15-Jun	2-Jul	59
2003	22-Jun	9-Jun	2-Jul	36
2004	15-Jun	5-Jun	30-Jun	57
2005	20-Jun	9-Jun	4-Jul	101
2006	29-Jun	19-Jun	12-Jul	79
2007	23-Jun	10-Jun	4-Jul	68
2008	23-Jun	13-Jun	10-Jul	73
Mean	26-Jun	17-Jun	8-Jul	38

Year	Average	Min	Max	N
Common Eider				
1982	9-Jul	8-Jul	10-Jul	4
1983	26-Jun	21-Jun	30-Jun	3
1984	--	--	--	--
1985	--	--	--	--
1986	--	--	--	--
1987	29-Jun	25-Jun	8-Jul	10
1988	--	--	--	--
1989	2-Jul	29-Jun	8-Jul	4
1990	22-Jun	21-Jun	24-Jun	3
1991	26-Jun	19-Jun	5-Jul	27
1992	2-Jul	26-Jun	6-Jul	12
1993	24-Jun	18-Jun	27-Jun	5
1994	24-Jun	16-Jun	4-Jul	9
1995	23-Jun	14-Jun	2-Jul	13
1996	19-Jun	10-Jun	2-Jul	14
1997	19-Jun	10-Jun	1-Jul	15
1998	28-Jun	20-Jun	4-Jul	18
1999	30-Jun	22-Jun	9-Jul	12
2000	29-Jun	24-Jun	5-Jul	23
2001	30-Jun	20-Jun	8-Jul	23
2002	24-Jun	15-Jun	30-Jun	17
2003	22-Jun	14-Jun	4-Jul	16
2004	17-Jun	6-Jun	26-Jun	18
2005	19-Jun	5-Jun	1-Jul	34
2006	1-Jul	24-Jun	11-Jul	52
2007	20-Jun	15-Jun	30-Jun	50
2008	24-Jun	16-Jun	2-Jul	34
Mean	26-Jun	19-Jun	3-Jul	18

Table 1. Estimated hatch date continued.

Year	Average	Min	Max	N
Pacific Loon, Red-throated Loon				
1982	8-Jul	3-Jul	24-Jul	25
1983	29-Jun	21-Jun	29-Jul	15
1984	2-Jul	26-Jun	8-Jul	5
1985	7-Jul	25-Jun	21-Jul	15
1986	5-Jul	26-Jun	25-Jul	37
1987	3-Jul	27-Jun	12-Jul	34
1988	27-Jun	16-Jun	5-Jul	5
1989	2-Jul	22-Jun	15-Jul	5
1990	1-Jul	25-Jun	9-Jul	11
1991	26-Jun	18-Jun	5-Jul	21
1992	5-Jul	29-Jun	18-Jul	12
1993	26-Jun	18-Jun	5-Jul	12
1994	24-Jun	19-Jun	29-Jun	6
1995	26-Jun	21-Jun	1-Jul	10
1996	22-Jun	15-Jun	1-Jul	9
1997	22-Jun	15-Jun	29-Jun	17
1998	1-Jul	20-Jun	14-Jul	37
1999	3-Jul	22-Jun	14-Jul	48
2000	30-Jun	15-Jun	9-Jul	40
2001	4-Jul	27-Jun	15-Jul	27
2002	25-Jun	12-Jun	3-Jul	42
2003	24-Jun	12-Jun	3-Jul	14
2004	23-Jun	13-Jun	30-Jun	10
2005	27-Jun	11-Jun	7-Jul	42
2006	2-Jul	27-Jun	7-Jul	22
2007	24-Jun	15-Jun	2-Jul	31
2008	28-Jun	16-Jun	9-Jul	46
Mean	29-Jun	20-Jun	10-Jul	22

Year	Average	Min	Max	N
Glaucous Gull				
1982	5-Jul	29-Jun	22-Jul	23
1983	22-Jun	13-Jun	4-Jul	14
1984	23-Jun	18-Jun	26-Jun	5
1985	3-Jul	23-Jun	12-Jul	23
1986	27-Jun	22-Jun	5-Jul	18
1987	28-Jun	20-Jun	10-Jul	19
1988	22-Jun	15-Jun	3-Jul	9
1989	22-Jun	22-Jun	22-Jun	3
1990	--	--	--	--
1991	18-Jun	12-Jun	3-Jul	26
1992	27-Jun	22-Jun	4-Jul	23
1993	20-Jun	15-Jun	7-Jul	11
1994	17-Jun	10-Jun	27-Jun	17
1995	17-Jun	14-Jun	26-Jun	17
1996	14-Jun	11-Jun	20-Jun	15
1997	17-Jun	10-Jun	29-Jun	19
1998	22-Jun	15-Jun	9-Jul	64
1999	27-Jun	19-Jun	7-Jul	25
2000	22-Jun	12-Jun	9-Jul	72
2001	24-Jun	17-Jun	7-Jul	50
2002	17-Jun	6-Jun	4-Jul	56
2003	13-Jun	4-Jun	26-Jun	58
2004	10-Jun	3-Jun	19-Jun	21
2005	14-Jun	6-Jun	27-Jun	69
2006	25-Jun	17-Jun	9-Jul	46
2007	15-Jun	8-Jun	29-Jun	76
2008	20-Jun	9-Jun	1-Jul	68
Mean	21-Jun	14-Jun	3-Jul	33

Year	Average	Min	Max	N
Mew Gull				
1982	10-Jul	7-Jul	22-Jul	11
1983	26-Jun	17-Jun	3-Jul	6
1984	--	--	--	--
1985	4-Jul	27-Jun	12-Jul	8
1986	2-Jul	21-Jun	12-Jul	18
1987	26-Jun	21-Jun	4-Jul	8
1988	18-Jun	14-Jun	24-Jun	4
1989	--	--	--	--
1990	--	--	--	--
1991	20-Jun	14-Jun	2-Jul	8
1992	27-Jun	23-Jun	4-Jul	10
1993	24-Jun	17-Jun	2-Jul	7
1994	15-Jun	11-Jun	21-Jun	8
1995	18-Jun	15-Jun	22-Jun	16
1996	14-Jun	8-Jun	20-Jun	10
1997	19-Jun	16-Jun	27-Jun	8
1998	24-Jun	19-Jun	4-Jul	19
1999	25-Jun	21-Jun	9-Jul	25
2000	25-Jun	17-Jun	5-Jul	17
2001	26-Jun	19-Jun	7-Jul	18
2002	16-Jun	6-Jun	3-Jul	40
2003	17-Jun	8-Jun	27-Jun	20
2004	13-Jun	9-Jun	19-Jun	19
2005	19-Jun	10-Jun	1-Jul	32
2006	26-Jun	18-Jun	9-Jul	45
2007	18-Jun	7-Jun	3-Jul	32
2008	19-Jun	13-Jun	4-Jul	45
Mean	23-Jun	16-Jun	2-Jul	18

Year	Average	Min	Max	N
Sabine's Gull				
1982	--	--	--	--
1983	21-Jun	14-Jun	2-Jul	3
1984	--	--	--	--
1985	2-Jul	26-Jun	18-Jul	3
1986	24-Jun	15-Jun	7-Jul	7
1987	21-Jun	15-Jun	4-Jul	7
1988	24-Jun	18-Jun	8-Jul	7
1989	--	--	--	--
1990	--	--	--	--
1991	15-Jun	9-Jun	22-Jun	9
1992	--	--	--	--
1993	17-Jun	14-Jun	23-Jun	8
1994	11-Jun	9-Jun	16-Jun	6
1995	18-Jun	12-Jun	28-Jun	6
1996	11-Jun	7-Jun	14-Jun	3
1997	14-Jun	8-Jun	22-Jun	8
1998	21-Jun	15-Jun	6-Jul	11
1999	21-Jun	16-Jun	3-Jul	20
2000	22-Jun	14-Jun	2-Jul	7
2001	27-Jun	19-Jun	4-Jul	10
2002	14-Jun	8-Jun	26-Jun	28
2003	12-Jun	6-Jun	17-Jun	5
2004	9-Jun	3-Jun	19-Jun	3
2005	16-Jun	8-Jun	29-Jun	30
2006	22-Jun	18-Jun	27-Jun	23
2007	17-Jun	9-Jun	29-Jun	30
2008	17-Jun	10-Jun	26-Jun	17
Mean	19-Jun	13-Jun	29-Jun	11

Table 1. Estimated hatch date continued.

Year	Average	Min	Max	N	Year	Average	Min	Max	N
Arctic Tern					Greater Scaup, Long-tailed Duck				
1982	--	--	--	--	1982	10-Jul	13-Jul	22-Jul	5
1983	--	--	--	--	1983	--	--	--	--
1984	--	--	--	--	1984	--	--	--	--
1985	29-Jun	22-Jun	4-Jul	8	1985	--	--	--	--
1986	26-Jun	16-Jun	24-Jul	6	1986	8-Jul	1-Jul	11-Jul	4
1987	24-Jun	20-Jun	26-Jun	3	1987	7-Jul	2-Jul	11-Jul	4
1988	--	--	--	--	1988	--	--	--	--
1989	--	--	--	--	1989	--	--	--	--
1990	--	--	--	--	1990	--	--	--	--
1991	17-Jun	12-Jun	20-Jun	4	1991	--	--	--	--
1992	1-Jul	25-Jun	10-Jul	6	1992	6-Jul	30-Jun	22-Jul	19
1993	17-Jun	15-Jun	20-Jun	3	1993	2-Jul	29-Jun	8-Jul	18
1994	--	--	--	--	1994	2-Jul	30-Jun	3-Jul	7
1995	16-Jun	13-Jun	20-Jun	3	1995	3-Jul	20-Jun	10-Jul	14
1996	--	--	--	--	1996	28-Jun	17-Jun	5-Jul	7
1997	--	--	--	--	1997	27-Jun	17-Jun	4-Jul	10
1998	26-Jun	19-Jun	5-Jul	5	1998	4-Jul	26-Jun	10-Jul	14
1999	25-Jun	21-Jun	2-Jul	8	1999	8-Jul	1-Jul	13-Jul	10
2000	26-Jun	23-Jun	1-Jul	5	2000	6-Jul	28-Jun	11-Jul	26
2001	22-Jun	15-Jun	29-Jun	5	2001	7-Jul	4-Jul	17-Jul	6
2002	18-Jun	8-Jun	26-Jun	37	2002	1-Jul	29-Jun	4-Jul	8
2003	13-Jun	8-Jun	21-Jun	5	2003	30-Jun	29-Jun	1-Jul	3
2004	17-Jun	9-Jun	29-Jun	9	2004	--	--	--	--
2005	21-Jun	14-Jun	29-Jun	15	2005	1-Jul	24-Jun	8-Jul	9
2006	23-Jun	17-Jun	29-Jun	17	2006	5-Jul	29-Jun	8-Jul	5
2007	16-Jun	8-Jun	24-Jun	18	2007	2-Jul	1-Jul	5-Jul	5
2008	20-Jun	11-Jun	30-Jun	16	2008	--	--	--	--
Mean	22-Jun	16-Jun	29-Jun	10	Mean	3-Jul	28-Jun	9-Jul	10
Pintail, Shoveler, Mallard, Teal					Small Shorebird				
1982	--	--	--	--	1982	--	--	--	--
1983	--	--	--	--	1983	--	--	--	--
1984	--	--	--	--	1984	--	--	--	--
1985	--	--	--	--	1985	3-Jul	22-Jun	17-Jul	4
1986	3-Jul	22-Jun	17-Jul	13	1986	27-Jun	16-Jun	11-Jul	23
1987	3-Jul	18-Jun	14-Jul	12	1987	21-Jun	15-Jun	25-Jun	7
1988	--	--	--	--	1988	--	--	--	--
1989	6-Jul	3-Jul	11-Jul	4	1989	24-Jun	18-Jun	4-Jul	3
1990	28-Jun	24-Jun	3-Jul	4	1990	--	--	--	--
1991	24-Jun	12-Jun	4-Jul	13	1991	22-Jun	8-Jun	6-Jul	21
1992	4-Jul	21-Jun	13-Jul	16	1992	26-Jun	22-Jun	29-Jun	8
1993	28-Jun	24-Jun	1-Jul	16	1993	21-Jun	14-Jun	29-Jun	15
1994	28-Jun	26-Jun	29-Jun	5	1994	--	--	--	--
1995	27-Jun	18-Jun	6-Jul	11	1995	19-Jun	10-Jun	25-Jun	6
1996	25-Jun	13-Jun	2-Jul	10	1996	20-Jun	8-Jun	2-Jul	10
1997	17-Jun	13-Jun	21-Jun	4	1997	12-Jun	9-Jun	14-Jun	3
1998	1-Jul	18-Jun	10-Jul	39	1998	22-Jun	14-Jun	28-Jun	8
1999	2-Jul	20-Jun	12-Jul	17	1999	25-Jun	16-Jun	5-Jul	17
2000	30-Jun	21-Jun	8-Jul	28	2000	23-Jun	14-Jun	27-Jun	13
2001	2-Jul	27-Jun	8-Jul	13	2001	27-Jun	18-Jun	4-Jul	22
2002	24-Jun	13-Jun	3-Jul	21	2002	19-Jun	12-Jun	25-Jun	21
2003	20-Jun	10-Jun	29-Jun	8	2003	16-Jun	13-Jun	17-Jun	3
2004	18-Jun	5-Jun	28-Jun	19	2004	12-Jun	8-Jun	20-Jun	10
2005	23-Jun	11-Jun	4-Jul	24	2005	20-Jun	9-Jun	29-Jun	30
2006	1-Jul	23-Jun	8-Jul	15	2006	27-Jun	19-Jun	8-Jul	29
2007	22-Jun	13-Jun	28-Jun	17	2007	18-Jun	5-Jun	28-Jun	33
2008	26-Jun	18-Jun	4-Jul	22	2008	19-Jun	11-Jun	1-Jul	54
Mean	27-Jun	18-Jun	5-Jul	15	Mean	21-Jun	13-Jun	30-Jun	16